

A. argenteus. *Cand. cat. hort. monsp. p. 77. Humb. Bonpl. Kunth nov. gen. et sp. 1. p. 188. Roem. Schult. syst. veg. 2. p. 815.*

Saccharum argenteum. *Brouss. el. p. 50. Jacq. ecl. gram. fasc. 1. t. 5.*

Erianthus saccharoides. *Willd. enum. 1. p. 46. excl. syn. et charact. gener.*

Hab. in montanis huanoccensibus Peruviae. 2.

Flos neuter uniglumis.

**** *Panicula.*

ANDROPOGON SUBULATUS. PRESL.

A. culmo basi repente, ramoso, nodisque vaginisque glabro, foliis abbreviatis, lanceolato-linearibus, margine scabris, paniculae coarctatae ramis simplicissimis verticillatis, rhachi ramisque scabris, calcare e ramis accedente hirsuto, locustis ternis, media sessili subbiflora, monica, gluma inferiore limucronata, superiore aristata, paleis ciliatis, inferiore feminei flosculi apice aristata, locustis lateralibus pedicellatis masculis unifloris, glumis setigeris, paleis ciliatis.

Hab. in Luzonia ad Sorzogon. 2.

Valde affinis *A. sciculari* Retz, sed omnino distinctus, praesertim, si comparamus iennem et descriptionem a cl. Trinio in speciebus graminum datam, ab hac non solum planta Haekiana, imo exemplaria mauritiana a cl. Sieber data discrepant. E descriptione nostra descriptio elucescent.

Radix repens. Culmus inferne repens, adscendens, erectus, compressus, alternatim hinc at illinc canaliculatus, in 1—2 ramos divisus, glaber, pedalis et altior. Folia infima vaginis longiora, margine hispidula, superiora vaginis multoties breviora, rudimentalia, acutiuscula. Vaginae glabrae. Ligula brevissima, ciliata. Panicula 3-pollicaris, lineari-lanceolata, simplicissima, contracta. Rhachis obtusangula, striata, scabra, partialis s. rami ad medietatem fere a superioribus longitudinaliter demum accedentes in partes duas, ramulum nimirum et calcar; ramulus persistens obverse setaeformis, apice acutus, teretiusculus, supra medium uno latere canaliculatus, quo recipitur calcar immediate locustas gerens, claviculatus, triquetrum, latere externo convexum, a basi debians et cum locustis decidens, hirsutum pilis sursum incrementibus et apice locustas obvallantibus, rufescentibus. Locustae ternae, sessilis et binae pedicellatae. Sessilia subbiflora, subulata. Glumae duae: inferior lanceolata, dorso convexa, apice fissa, longe bimucronata, cartilaginea, margine inflexa, paleam flosculi feminei superiorem ambiens, obsolete binervia, nervis mucronibusque hispidis; superior lanceolata, carinata, uninervia, nervo supra medium margineque hispida, apice bidenticulata, inter dentes aristam dimidio brevioram scabram rectam setaeformem emittens. Flosculi duo, uno femineo inferiore altero masculo superiore saepissime abortivo. Flosculus femineus bipaleaceus: palea inferior lanceolato-linearis, glumae inferiori opposita et superiori opposita, his duplo brevior margine sat dense ciliata, diaphana, mollis, apice in aristam fere triplo longiorem rectam scabram excurrentem, saepe gluma superiore recepta; palea superior glumae inferiori apposita, inferiore duplo longior et latior, paulo brevior gluma inferiore, lanceolata, diaphana, integerrima, margine ciliata. Lodica foliolis duobus ovatis truncatis. Ovarium lineari-lanceolatum, lodicula brevius. Styli duo. Stigmata aspergilliformia. Caryopsis lineari-lanceolata, apice mucronata, mucrone pubescente, rufa, libera. Flosculus superior uniglumis: gluma superior ovata, binervia, bicarinata, carinis s. nervis scabra, integerrima, inferiora vix supplet palea superior; stamen unicum; flosculus hic saepissime abortivus, palea staminisque deficientibus. Locustae pedicellatae masculae uniflorae subulatae, sessilem longitudine aequantes. Pedicelli locustae sessili vix breviores, semiteretes, latere interno concavusculi canaliculati, sursum parum incrassati, apice truncati, scabriusculi, apice pilis rigidis sursum vergentibus subadpressis hispiduli. Glumae dorso convexae, purpureo virscentes, inferior lanceolata, trinervia, scabriuscula, apice subulato elongata, bidenticulata et in aristam duplo brevioram excurrentem, subula et arista scabra, superior conformis, paulo longior, marginibus ciliata, membranacea, 3-nervia apice in subulam et aristam breviores producta, includens flosculum. Palea inferior paulo brevior gluma lanceolata, diaphana, 3-nervia, acutissima, margine dense piloso-ciliata, superior tertia parte brevior, lanceolata, integerrima, margine ciliata. Stamina tria.