

Genus 2, Telea, Hübner.

1. *T. Polyphemus*, *Fab.*—Abundant, June and July.

Distribution.—Brazil, (B.M.C.); Massachusetts, (Harris); New York, (Emmons); Eastern Township (Gosse); Rouge District.

Genus 3, Tropœa, Hübner.

1. *T. Luna*, *Linn.*—Rare, June.

Distribution.—Mexico, (B.M.C.); Massachusetts, (Harris); New York, (Emmons); Niagara, Sorel, &c.

*Group 3. NOCTUINA, Stainton.**Division 1. TRIFIDÆ, Guénée.**Sub-Division 1. BOMBYCIFORMES, Guénée.**Family. CYMATOPHORIDÆ, Hen.-Schæff.**Genus 1, Gonophora, Bruand; Thyatira, Ochs., &c., p.*

1. *G. scripta*, *Gosse.*—Not very common, June and July.

Distribution.—New York, (E. Doubleday); Eastern Townships, (Gosse); Orillia, C.W., (Bush); Rouge District.

Genus 2, Thyatira, Ochs.

1. *T. cymatophoroides*, *Guén.*—Common, June and July.

Distribution.—New York, (E. Doubleday); Orillia, C.W., (Bush); Rouge District.

Genus 3, Leptina, Guén.

1. *L. Doubledayi*, *Guén.*—Rare.

Distribution.—Northern States, (E. Doubleday).

*FAMILY, BOMBYCOIDÆ, Guén.**Genus, 1. Balsa, Walker M.S.S.*

"*Balsa*, N.G. *Mas.* Corpus gracile. Proboscis distincta. Palpi graciles, recti, subpilosi, oblique ascendentes; articulus 2 us. longus; 3 us. parvus, lanceolatus. Antennæ glabræ. Abdomen subcarinatum, apice compressum, alas posticas non superans. Pedes graciles, glabri; calcaria longa. Alæ latiusculæ, apice subrectangulatæ, costa vix convexa, margine exteriori subconvexa subobliquo."

"*Male.* Body slender. Proboscis distinct. Palpi slender, straight, slightly pilose, obliquely ascending a little higher than the head: second joint long: third minute, lanceolate. Antennæ simple, smooth. Abdomen slightly keeled not extending beyond the hinder wings; tip compressed, with a small apical tuft. Legs slender, smooth; hind tibiæ with four long spurs. Wings rather broad. Forewings nearly rectangular at the tips; exterior border slightly convex and oblique; costa very slightly convex."