

male palpi are quite simple in structure. Our genera may be separated by the following table :

1	{	M. E. *in front of A. S. E.....	<i>Usofila</i>
		M. E. not as far front as A., 'S. E.....	2
2	{	M. E. in front of P. S. E.....	<i>Segestria</i>
		M. E. as far back as P. S. E.....	3
3	{	Three claws to tarsus.....	<i>Ariadne</i>
		Two claws to tarsus.....	<i>Dysdera</i>

USOFILA GRACILIS, Keys.

Marx in Proc. Ent. Soc., Wash., 1890, Vol., II., p. 36, pl. I., fig. 6.
Alabaster Cave, Cala.

This is quite unlike other *Dysderidæ* in general appearance, but is referred by Keyserling to this family.

DYSDERA INTERRITA, Hentz.

D. crocata, Koch. Marx Cat. Aranæ of temp. of N. Am., 1890.

D. interrta, Hentz. Proc. Bost. Soc. Nat. Hist., Vol. IV., p. 224, 1842.

“ “ “ Spid. U. S., ed. Burgess, p. 20, pl. II., fig. I., 1875.

“ “ “ Emerton, New Engl. Drass., Agal. and Dysd.,
p. 36, pl. VIII., fig. 2, 1890.

New Engl., N. Y., D. C., Md., Va.

ARIADNE BICOLOR, Hentz.

Pylarus bicolor, Hentz. Proc. Bost. Soc. Nat. Hist., Vol. IV, p. 225,
1842.

“ “ “ Spid. U. S., ed. Burgess, p. 21, pl. II., fig.
3, 1875.

“ *pumilis*, Hentz. Proc. Bost. Soc. Nat. Hist., Vol. IV., p.
226, 1842.

“ “ “ Spid. U. S., ed., Burgess, p. 22, pl. II.,
fig. 5, 1875.

*M.E.—median eyes; A.S.E.—anterior side eyes; P.S.E.—posterior side eyes,