summer, and thus betrays its presence. Time of stay: 6th of May to 22nd of September.

197. Lanivireo flavifrons, Yellow-throated Vireo. A rare summer resident. In six years I have seen four here, two of which were on King's Mountain, July 20th, 1905. They arrive about May 11th.

198. Lanivireo solitarius, Blue-headed Vireo. A moderately common migrat, but rare breeder. They arrive May 7th, and the latest date in them is September 11th (1909). On May 28th 1898, a nest was found at Chelsea, and on August 21st, 1907, I saw fully grown young on Kettle Island.

MNIOTILTIDÆ-WOOD WARBLERS.

199. Mniotilta varia, Black and White Warbler. A common migrant and moderately common breeder. This frequenter of the more open, deciduous second-growth woods arrives from April 28th to May 13th; the last are seen September 21st. Breeds in Beaver Meadow, Blueberry Point and similar places.

200. Vermivora rubricapilla. Nashville Warbler. A moderately common migrant and breeder, but only in its chosen haunts, viz., the stands of small poplar and birch in low, wet places in woods. In such a spot at Blueberry Point a pair or two can be found any day in summer. More common in the poplar stands around the Mer Bleue. They begin to arrive May 7th, reaching their climax in numbers May 15th. On July 13th, 1881, a nest with four eggs was found at Dow's Swamp. The latest date for them is September 25th.

201. Vermivora celata, Orange-crowned Warbler. An extremely rare migrant. On September 27th, 1885, one was taken by Mr. E. White near the eastern end of the city. This is the only record.

202. Vermivora peregrina, Tennessee Warbler. A rare migrant; will probably be found breeding in some of the tamarack swamps in the northern part of the district. It passes through Ottawa from May 12th to June 7th (1907), at which latter date Messrs. Chapman and Fuertes saw some in Major's Hill Park. The last are seen September 30th. Its song is a high, sharp tsip, tsip, tsip, pit it it itereeeee.

203. Compsothlypis americana usneæ, Northern Parula Warbler. A moderately common migrant and rarer breeder. Its habitat is tall deciduous trees, as on the west side of Beaver Meadow, where it may be found in summer. It begins to arrive May 7th, reaching its climax May 20th to 23rd. The last are seen September 13th. The song is a somewhat scratchy, dreer, dreer, dreer, last note highest and loudest; also a trill like re-e-e-e-e-e.

(To be continued.)