

followed by the same condition. At period of examination she complained of a feeling of impending dread, and manifested suicidal impulses.

Pelvic examination showed retroversion with adhesions.

Treatment recommended.

No. 139.—Mrs. H., aged 36. One brother slightly deranged. Has had three children, followed by several miscarriages. After the last miscarriage she had "fever," which, according to her husband, left her insane, necessitating her confinement to an asylum for three months, afterwards returning to her home, but unfit to properly manage it. Became suspicious of friends, abusive to the children and wholly unreliable.

Examination showed enlarged uterus and deep cervical tear. Amputation of cervix, removal of uterine fungosities, resection of the tubes resulted in but slight mental improvement. Patient has passed from observation.

No. 140.—Mrs. P., aged 49. Heredity good. Dementia of seven years' standing with periods of deep melancholia.

Examination showed right floating kidney and retroversion. Fixation of kidney and suspension of uterus done, no appreciable result until some three months afterwards when her husband wrote:—"Mrs. P. is recovering fast, both in body and mind, she is around and doing the most of her work. I feel so thankful that she is getting all right again."

No. 141.—Mrs. W., aged 65. Good heredity. Religious mania with melancholy lasting several months. Several years previously she had an ovarian cyst removed. She complained of pain in left inguinal region. Pelvic examination was negative. Upon opening the abdomen the omentum was found adherent to the abdominal wall, and the rectum adherent to the side of the uterus and to the stump of the tube. These adhesions were carefully freed.

Death followed from obstruction of the bowels. A post-mortem examination showed a knuckle of ilium had become adherent within the pelvis.

No. 142.—Mrs. D., aged 44. Referred to by Dr. Jeffs. Good heredity. No miscarriages. Youngest child 7 years old. Mild religious delusions for several years. For two weeks were so decided that removal to an asylum was contemplated.

Deep laceration of cervix, small cyst of left ovary, myometritis with fungosities. The appropriate treatment was given. Gradual return to mental health, after two months Dr. Jeffs reports her as recovered.

No. 143.—Mrs. W., aged 36, three children, youngest 10 years,