tergite broad and long with a cross-shaped mark; near its tip—set with two small, semicircular, chitinized pieces which are produced into sharp points on the proximal ends. Pleurites short and stout, at the base on the dorsal side, produced into a long, slender, chitinized arm which is directed entad, two apical appendages, the more ventrad being chitinized, especially at the tips, the dorsal apical appendage fleshy. Between the tergite and the unarmed sternite, nearly in the median plate, is a rectangular, subchitinized organ, bearing at its outer angles chitinized hooks, bent ventrad and inward, these hooks minutely denticulated at tip.

J.-Length, 5.8 mm.; wing, 6.3 mm.

♀.—Length, 6.4-7.1 mm.

Holotype.-Vial 6, April 25, 1912; Tokio, Japan.

Allotype.-Vial 6, April 25, 1912; Tokio, Japan.

Paratypes.—Vial 6 and L; 4  $\, \, \, \, \diamondsuit$  , 2  $\, \, \, \, \circlearrowleft$  , April 25, 1912; Aug. 1912, Tokio, Japan.

Subgenus Erioptera, Meigen.

Erioptera (Erioptera) elegantula, sp. n.

Wings with brown spots.

Male.-Length, 5.4 mm.; wing, 7.7-7.9 mm.

Female.-Length, 6-6.5 mm.; wing, 7-8.3 mm.

Male.—Rostrum and palpi dark brown, antennæ with basal segments brown, flagellar segments short, dark brown; front, ver-

tex and occiput dark brown.

Pronotum dark brown above, lighter colored on the sides. Mesonotum dark brown, the region before the pseudosutural pits more yellowish; scutum, scutellum and postnotum dark brown. Pleuræ dark brown. Halteres pale. Legs: coxæ dark brown; trochanters brown; femora dark brown; tibiæ dark brown, a little paler at the extreme base; tarsi dark brown. Wings subhyaline with greyish-brown marks, as follows: A large rounded spot at origin of Rs, a second at Sc², a third at end of Sc¹ running down over cross-vein r; a fourth spot at tip of R¹ and a smaller one at tip of R²; cord broadly margined with the same color; less distinct clouds at ends of the other veins and along most of these veins. Venation, (see fig. 3, plate III.)

Abdomen dark brown, densely clothed with long whitish hairs. Hypopygium. 9th tergite broad at base, narrowed at the middle,