

continuation on the mucous membrane, is the only mark of the operation: it fairly halves the lip, and the two portions are perfectly symmetrical. Instead of the centre of the prolabium being notched, it is rather dependant, as in lips supposed, physiognomically, to indicate the faculty of concentration. Although union seemed firm, straps, &c., were re-applied by way of precaution, and advised to be kept on a few days longer. He left the Hospital this afternoon to join his ship at Quebec, as she was to leave on her homeward voyage the next day.

The subjoined cuts, taken from drawings made by Dr. A. C. Mac-Donell, are faithful representations of the lip before and after the operation:—



RETURN of Sick in the Marine and Emigrant Hospital, Quebec, from the 31st July to the 3rd September, 1853, inclusive.

	Men.	Women.	Children.	Total.
Remained, Since admitted,	61	15	2	78
	120	34	10	164
	181	49	12	242
Discharged,	117	22	4	141
Died,	7	3	1	11
Remaining,	57	24	7	88
	181	49	12	242

Fever,	72	Amenorrhœa,	1
Inflammation of Lungs,	5	Rubeola,	2
Inflammation of Bowels,	1	Febricula,	7
Dyspepsia,	2	Disease of Knee Joint,	2
Rheumatism,	6	Pregnancy,	1
Dysentery,	10	Albuminuria,	1
Small Pox,	1	Ophthalmia,	2
Dropsy,	1	Subluxatio,	2
Diseases of Skin,	2	Feb. Intermittent,	1
Syphilis,	12	Scorbutus,	4
Fractures,	3	Strictures,	1
Abscess,	3	Necrosis os Maxil,	1
Ulcers,	4	Delirium Tremens,	1
Wounds,	2	Insanity,	1
Contusions,	11	Paralysis,	1
Burns and Scalds,	1		

C. E. LEMIEUX, House Surgeon.