

AND SIMILAR PREPARATIONS—*Concluded.*

Directions for use, as found on the Label.	RESULTS OF ANALYSIS.						Remarks.	
	Weight of Contents of one Powder.		Acetani- lide= anti- febrin.	Phen- acetin.	Phena- zone= anti- pyrine.	Caffeine		Car- bonate of Soda.
	Grms.	Grms.						
Directions—Place a powder on the tongue, swallow with a draught of water, if not relieved in half an hour take another powder. Do not repeat again for 3 or 4 hours. For children $\frac{1}{4}$ to $\frac{1}{2}$ a powder. E. A. Goodman, chemist, 380 Yonge street, Toronto.	9.0	0.58	Present.	None ..	None ..	Doubt- ful.	Present.	
These tablets immediately relieve nervous and sick headache, neuralgia, pains in any part of the body; perfectly safe—do not contain any opiates—are new and very efficacious. Dose—Two tablets at once; repeat by taking 1 in $\frac{1}{2}$ hour if necessary. Pour les directions en Francais voir le circulaire inclus.	6.5	0.42	" ..	" ..	" ..	Present.	" ..	
Directions—Place this powder on the tongue and swallow with a draught of water, or stir the powder briskly in water and drink quickly; if not relieved in one hour take another powder; children under twelve years old, half a powder.	12.0	0.78	" ..	" ..	" ..	Doubt- ful.	None ..	