

NYCTEA NYCTEA. Snowy Owl.

SURNIA U. CAPAROCH. American Hawk Owl. Rare.

GLAUCIDIUM GNOMA CALIFORNICUM. Pigmy Owl. Common.

CEKYLE ALCYON. Belted King-fisher. Quite a number stay all winter.

CEOPHLEUS PILEATUS. Pileated Wood-pecker. Tolerably common.

DRYOBATES V. LEUCOMELAS. Northern Hairy Woodpecker.

" P. ORCUEUS. Batchelder's Woodpecker.

Both common.

XENOPICUS ALBOLARVATUS. White Headed Woodpecker. I heard of this species but never came across it.

PICOIDES ARCTICUS. Arctic 3-toed Woodpecker. Common in suitable localities.

COLAPTES CAFER. Red-shafted Flicker. Fairly common during winter months.

OTOCORIS SP? Horned Lark. On January 10th, '98, I saw a flock of shorelarks, most probably *merrilli*, but they may have been *strigata* as both species occur and are common, the latter only on migrations, but the former breeds.

PICA HUDSONICA. American Magpie. Abundant.

CYANOCETTA S. ANNECTENS. Black Headed Jay. Common.

PERISOREUS C. CAPITALIS. Rocky Mt. Jay. Common at high elevations.

CORVUS C. PRINCIPALIS. Northern Raven. Tolerably common.

CORVUS AMERICANUS HESPERIS. California Crow. A few crows stay around the towns, but the bulk leave rather early in the fall.

PICICORVUS COLUMBIANUS. Clarke's Nutcracker. Common during winter of '97-'98, but entirely absent the next winter. Many remained to breed in '98 and they also bred in '97; they lay in February.

AGELAI VS PH. SONORIENSIS. Sonoran Redwing.

SCOLECOPHAGUS CYANOCEPHALUS. Brewer's Blackbird.

A few of both the above species of Blackbirds remain throughout the winter.

STURNELLA NEGLECTA. Western Meadow Lark. A number remain about the stacks and barnyards.

COCCOTHRAUSTES VESP. MONTANUS. Western Evening Grosbeak. Tolerably common.

PINICOLA E. CANADENSIS. Pine Grosbeak. Scarce during winter of '97-'98, but common during '98-'99. Breeds.

LOXIA C. MINOR. American Crossbill. Abundant during winter of '97-'98, but apparently entirely absent the next season. I never positively identified the white-winged Crossbill, but I have no doubt it occurs as I have taken it west of the Cascades.

LEUCOSTICTE TEPHIROCOTIS. Gray-crowned Leucosticte. A large flock seen once or twice in mountains to west of Okanagan Lake; these may have been *littoralis* as I have taken both species in lower Fraser River Valley.

ACANTHIS LINARIA. Redpoll. Common during both winters I spent in the district. I saw nothing of *exilipes*, though I kept a good look out for it and have taken it nearer the coast.

SPINUS TRISTIS. American Goldfinch. Common during early part of winter of '97-'98; never seen since. This is probably the western form.

SPINUS PINUS. Pine Finch. Abundant.