

*Andrena saccata*, n. sp.

♀ 13 mm. Face with black pubescence, dorsulum with pale pubescence, abdomen with erect, black pubescence.

Type locality: Corvallis, Or. Type Acad. Nat. Sci., Phila.

♀, 3rd May, 1899; 15th May, 1897; 8th June, 1898, Corvallis, Or.; 17th April, 1896, Seattle, Wash. (T. Kincaid). ♂, no data.

*Andrena hemileuca*, n. sp.

♀ 10 mm.

Type locality: Washington. Type Am. Ent. Soc., Phila.

♀, Seattle, Washington; Mt. Hood, Or.; ♂, Washington.

*Andrena clypeoporaria*, n. sp.

♀ 11 mm.

Type locality: Olympia, Wash. Type Acad. Nat. Sci., Phila.

♀, 12th June, 1895, Olympia, Wash.; Mt. Hood, Or.

*Andrena advarians*, n. sp.

♀ 13 mm.

Type locality: Vancouver, British Columbia. Type Acad. Nat. Sci., Phila.

♀, 5th April, 1902 (Harvey); ♂, 2nd March, 1902 (Harvey).

*Andrena Washingtoni*, Ckll., Psyche, IX., 1901, p. 284.

♀, 2nd June, 1895, Olympia, Wash. (Kincaid); 20th May, 1896, Seattle, Wash.

♂, 4th April, 1896, Seattle, Wash.

*Andrena moesta*, Sm. New Spec. Hym. Brit. Mus., p. 54.

♀, 6th May, Corvallis, Or. (Cordley).

*Andrena albihirta*, Ashm., ♀ = *perarmata*, Ckll., ♂. Bull. Col. Ass., I., p. 5.

♀, 13th, 17th March, 1896, Seattle, Wash. (T. Kincaid); Pullman, Wash. (C. V. Piper); 20th April, 1896, Livingston, Vanc.; 9th June, 1903, Vancouver, B. C. (Harvey).

♂, 16th, 18th February, 13th March, 1896; 15th March, 1897, Seattle, Wash. (T. Kincaid).

*Andrena Harveyi*, n. sp.

♀ 9 mm. Related to *A. mandibularis*, Rob.

Type locality: Grouse Mt., Vancouver, B. C.

Type Acad. Nat. Sci., Phila.