

Respiratory system.—Empyema	2
Pharyngitis, etc	8
Pleuritis	13
Pneumonia	8
Osseous muscular system, etc.—Arthritis	1
Mastoiditis	1
Myositis	2
Osteomyelitis	5
Periostitis	4
Generative system.—Abortion	2
Mastitis	2
Urinary System.—Cystitis	9
Pyelitis	3
Nervous System.—*Paralysis	2
Post-typh. insanity	5
Tetany	2
Circulatory system.—Dilatation of heart	14
Pericarditis	1

The number of cases noted to have had cardiac dilatation is doubtless far from the true number, but slight differences of size of the heart, and transitory murmurs, indicating temporary dilatations are uncertain signs, and in the hands of house physicians of comparatively short experience are apt to be overlooked; so also many passing conditions of myocarditis of slight degree must escape notice.

ASSOCIATED CONDITIONS.—

Alcoholism	1	Orchitis	1
Arterio-sclerosis	1	Parasites	..
Cardiac valve lesions	16	Ascaris	5
Chlorosis	3	Taenia	1
Diphtheria	1	†Pregnancy	5
Epilepsy	2	Rheumatism, acute	2
Gonorrhoea	4	Syphilis	3
Hysteria	4	Tuberculosis pul	6
Neuritis, alcoholic	1		

HÆMORRHAGE: Hæmorrhage occurred in 72 cases, of which 26 proved fatal; in 11 cases hæmorrhage and perforation were associated. In 31 of these 72 cases, there are notes of the time at which food, other than purely liquid, was given; of these 31 cases, 12 had such food before hæmorrhage occurred. Although the notes are not absolutely explicit, it may be presumed that such diet was continued up to the time at which hæmorrhage occurred, and the space of time between the administration of such food and the occurrence of hæmorrhage varied from a few hours up to 29 days; the exact figures are as follows: immediately, 3 days, 5, 6, 7, 9, 9, 10, 11, 12, 12, 13, 26, 29.

* One case had the muscles of the face partially paralysed, the other the muscles of deglutition.

† In two cases abortion at 1st and 2nd month occurred; pregnancies of 5, 6½ and 7 months were uninterrupted.