

1842.

MAY, begins on SUNDAY.

Last Quarter, 2d. d. 8h. 32m. morning, ☾ bearing S. W.
 New Moon, 10th. d. 7h. 24m. morning, ☾ bearing E.
 First Quarter, 17th. d. 7h. 56m. morning, ☾ under the horizon.
 Full Moon, 24th. d. 5h. 25m. morning, ☾ under the horizon.

D	M	W	☾supper limb.		Days length.	☾fast of clock.	☾ dec. N.	☾ rises	☾ pl.	☾ south- ing.	Days incr's.
			rises.	sets.							
1	Su		4 49	7 5	14 15	3 2	15° 2'	m. 44	☾	5 24m	5 26
2	M		4 47	7 6	14 18	3 10	15 20	1 12	☾	6 10	5 29
3	Tu		4 46	7 8	14 21	3 17	15 38	1 36	☾	6 53	5 32
4	W		4 44	7 9	14 24	3 23	15 56	1 57	☾	7 34	5 35
5	Th		4 43	7 10	14 27	3 29	16 13	2 16	☾	8 14	5 38
6	F		4 42	7 11	14 29	3 34	16 30	2 35	☾	8 55	5 40
7	S		4 41	7 13	14 32	3 39	16 47	2 55	☾	9 37	5 43
8	Su		4 40	7 14	14 34	3 43	17 3	3 18	☾	10 21	5 45
9	M		4 38	7 15	14 37	3 46	17 19	3 46	☾	11 9	5 48
10	Tu		4 37	7 16	14 39	3 49	17 35	sets	☾	a. 1	5 50
11	W		4 36	7 17	14 41	3 51	17 51	8 55a.	☾	0 56	5 52
12	Th		4 35	7 18	14 43	3 53	18 6	10 4	☾	1 54	5 54
13	F		4 34	7 19	14 45	3 54	18 21	10 58	☾	2 54	5 56
14	S		4 33	7 20	14 47	3 55	18 36	11 40	☾	3 52	5 58
15	Su		4 32	7 21	14 49	3 55	18 50	morn.	☾	4 48	6 0
16	M		4 31	7 22	14 51	3 54	19 4	0 14	☾	5 41	6 2
17	Tu		4 30	7 23	14 53	3 53	19 18	0 42	☾	6 32	6 4
18	W		4 29	7 24	14 55	3 52	19 31	1 8	☾	7 21	6 6
19	Th		4 28	7 25	14 57	3 50	19 45	1 31	☾	8 10	6 8
20	F		4 27	7 26	14 59	3 47	19 57	1 55	☾	9 0	6 10
21	S		4 26	7 27	15 1	3 44	20 10	2 20	☾	9 51	6 12
22	Su		4 25	7 28	15 3	3 40	20 22	rises.	☾	10 44	6 14
23	M		4 24	7 29	15 5	3 36	20 33	7 5a.	☾	11 40	6 16
24	Tu		4 24	7 30	15 6	3 31	20 45	8 21	☾	8	6 17
25	W		4 23	7 31	15 8	3 26	20 56	9 20	☾	m 36	6 19
26	Th		4 22	7 32	15 10	3 20	21 6	10 5	☾	1 32	6 21
27	F		4 21	7 33	15 12	3 14	21 17	10 43	☾	2 26	6 23
28	S		4 20	7 34	15 14	3 7	21 27	11 14	☾	3 16	6 25
29	Su		4 19	7 35	15 16	3 0	21 36	11 39	☾	4 4	6 27
30	M		4 19	7 36	15 17	2 52	21 45	morn.	☾	4 48	6 29
31	Tu		4 18	7 37	15 19	2 44	21 54	0 1	☾	5 29	6 30

In light lands Buckwheat may be raised to advantage. It should not be sown till after the middle of May. One bushel is seed enough for an acre, if sown broad-cast, as is usual; but if sown in drills, less than half that quantity is sufficient. Buckwheat is harvested by mowing. After it is mown, it should be several days before it is housed. It is in no danger of the seeds falling, nor does it suffer much by wet. From its great succulency it is liable to heat in a mow, on which account it is better to put in into small stacks of five or six loads each, than either a large one, or in a barn. The most common mode of sowing