

I propose to state some additional facts respecting the species already known, to discuss their affinities, and to describe two additional species, making six in all from the Paleozoic rocks, including one from the Erian or Devonian. For reasons to be mentioned in the sequel, I do not admit the genus *Palæorbis* founded, by some German naturalists, on fossils which I believe to be tubes of Annelids.

It may be useful to premise that of the two leading subdivisions of the group of Pulmonifera, the Operculate and Inoperculate, the first has been traced no farther back than the Eocene. The second, or Inoperculate division, includes some genera that are aquatic and some that are terrestrial. Of the aquatic genera no representatives are known in formations older than the Wealden and Purbeck, and these only in Europe. The terrestrial group or the family of the *Helicidæ*, which, singularly enough, is that which diverges farthest from the ordinary gill-bearing Gasteropods, is the one which has been traced farthest back, and includes the Paleozoic species. It is further remarkable that a very great gap exists in the geological history of this family. No species are known between the Carboniferous and the early Tertiary, though in the intervening formations there are many fresh-water and estuarine deposits in which such remains might be expected to occur. There is perhaps no reason to doubt the continuance of the *Helicidæ* through this long portion of geological time, though it is probable that during the interval the family did not increase much in the number of its species, more especially as it seems certain that it has its culmination in the modern period, when it is represented by very many and large species, which are dispersed over nearly all parts of our continents.

The mode of occurrence of the Paleozoic Pulmonifera in the few localities where they have been found is characteristic. The earliest known species, *Pupa vetusta*, was found by Sir Charles Lyell and the writer, in the material filling the once hollow stem of a *Sigillaria* at the South Joggins in Nova Scotia, and many additional specimens have subsequently been obtained from similar repositories in the same locality, where they are associated with bones of Batrachians and remains of *Millipedes*. Other specimens, and also the species *Zonites priscus*, have been found in a thin, shaly layer, containing debris of plants and crusts of Cyprids, and which was probably deposited at the outlet of a small stream flowing through the coal-formation forest. The two species found in Illinois occur, according to Bradley, in an underclay or fossil soil which may have been the bed of a pond or estuary, and subsequently became a forest sub-soil. The Erian species occurs in shales charged with remains of land plants, and which must consequently have

receive  
in such  
pected  
Pulmon  
should  
minous  
Cyprid  
fossil p

With  
present  
sembla  
belong  
to sub  
nent a  
ble fro  
though  
small

All the  
moder

I sh  
of the  
known  
explor  
of the  
of exc  
anima  
ment  
the R  
three

[Sir  
the Sou  
1832 (6  
Air-bre  
384, 18

De

at th  
eter i  
adult  
ter o  
not  
teeth  
ing,  
spac  
in w  
spec  
rou