

# The Canadian Entomologist.

VOL. XXIV.

LONDON, MAY, 1892.

No. 5.

## MISCELLANEOUS NOTES ON BUTTERFLIES, LARVÆ, ETC.

BY W. H. EDWARDS, COALBURGH, WEST VA.

(Continued from page 56.)

### 2. *Argynnis Oweni*, n. sp.

MALE—Expands 2.25 inches.

Upper dull side red-fulvous, not much obscured by brown at base ; the markings rather heavy, the two marginal lines on both wings more or less confluent. Under side of primaries either yellow-buff from base to the row of rounded spots, the nervules being red-brown, or the interspaces are very lightly washed with this colour over the buff ground ; the subapical patch dark brown with a small imperfectly silvered spot ; the five submarginal spots also are of same character ; sometimes the silver is wholly wanting.

Secondaries dark brown and gray-buff, the brown prevailing on the disk, so that the second row of spots stand on a clear brown ground ; but above this to base the buff prevails, especially along the third row of spots, and below the cell ; but occasionally the mottled surface extends quite to the belt ; the belt red-buff, and very narrow, being much encroached on by the dark brown (almost black-brown) projections from the discal area, and by the wide brown edging of the submarginal spots ; these spots are small, semi-oval ; those of the second row small, sub-oval ; the outer row is generally well silvered, but all the others are imperfectly so in most cases, but occasionally one is well silvered throughout, or the outer two rows may be well and the remainder imperfectly silvered.

FEMALE.—Expands 2.4 inches.

The fulvous mottled with yellowish on both wings outside the mesial band, on secondaries the yellow corresponding with the second row of silver spots ; also in the cell of primaries the hue is yellow fulvous ; the black markings all very heavy ; the marginal lines nearly confluent throughout. Secondaries as in the male, except that the basal colour is