

neck white; the breast and belly grey and black mixed, interspersed with a few white spots; lower belly and thighs whitish, spotted with black; quills black, the first five marked with a white spot; the two middle feathers of the tail grey, crossed with five or six blackish bands; the others black, bordered with white, which takes up most space on the outer feathers; legs yellow brown.

Inhabits Cayenne, where it is found in the plantations; frequently quivers the wings, and utters a weak cry, like that of a toad; also another noise, not unlike that of a dog; it is not very shy, for it will suffer any one to come very near before it will fly away, and when disturbed, perches at no very great distance: it is not a rare species.

The tail in the *Pl. enlum.* is more than four inches long, and nearly even at the end; the middle feathers cinereous, marbled with black dots, with five or six narrow curved black bars; the exterior feather white, crossed about the middle with an oblique narrow black bar, the same near the tip; the two next feathers white within, but divided in the middle of the white with black, and a broad margin of black on the outer web, continuing round and including the end of the feather.

Only seen in Paraguay from September to November, hence may be supposed a Bird of Passage there; said have a note or cry similar to the word Chuyguignigui.

32. — GUIANA GOATSUCKER.

Caprimulgus Guianensis, *Ind. Orn.* ii. 586. *Gm. Lin.* i. 1030.

Le Montvroyan de la Guiane, *Bufl.* vi. 549.

Crapaud volant, ou Tette-chevre roux de la Guiane, *Pl. enl.* 733.

L'Ibityau jaspé, *Voy. d'Azara*, iv. No. 313.

Guiana Goatsucker, *Gen. Syn.* iv. 598. *Shaw's Zool.* x. 148.

LENGTH nine inches. Bill black, three quarters of an inch long, and beset with bristles; the general colour of the plumage fulvous, irregularly mixed with rufous, being in longitudinal streaks