are least cloathed with leaves; and is of a large size, being twelve inches deep, and sive or more in diameter, wholly composed of moss: this is only open on the side, about three inches from the top. The note is a soft whistle.

Le grand Gobe-mouche noir de Cayenne, Brif. orn. ii. p. 386. 15. pl. 38. f. 3.—Pl. enl. 381.

Le Piauhau, Buf. oif. iv. p. 588.

Lev. Muf.

SIZE of a Blackbird: length twelve inches. The bill is broad at the base, an inch and a quarter long, a little bent at the tip, and beset at the base with a sew hairs; the colour of it dusky: the whole plumage is black, except the chin,

throat, and fore part of the neck, on which is a large bed of beautiful crimson, inclining to purple: the legs are black.

Some specimens are wholly black, having not the least trace of crimson on the throat; and such may be esteemed as semales or young birds. I have seen more than one of the last, in which the plumage otherwise coincided exactly with the first-described.

These inhabit Cayenne, and other parts of South America: they are found in flocks, and precede in general the Toucans in their movements: feed on fruits and insects, to which last their bill seems fully adapted: are lively birds, always in action. Inhabit, for the most part, the woods, like the Toucans; and where the first are found, the others are feldom far off. They have a sharp cry, which they often repeat, not unlike Pibauhau; whence their name.

Linnæus

78. + PURPLE-THROATED FL.

DESCRIPTION.

VARIETIES.

PLACE AND MANNERS.