

parts of the country can doubt. Not only is *Taraxacum* not rare on the banks of lakes and streams, but the writer has often found it in bogs and swamps several hundred miles from settlement of any kind.

TARAXACUM CHAMISSONIS, Greene, Pittonia, vol. iv, p. 228.

Very common on the shores and islands of Behring Sea and south along the Alaskan coast. Will probably be found in British Columbia.

TARAXACUM RUPESTRE, Greene, Pittonia, vol. iv, p. 229.

Crevice of rocks, alt. 6,000 ft., Mt. Queest, Shuswap Lake, B. C. Herb. No. 15,111; Avalanche Mt., Selkirk Mountains, B. C., alt. 8,000 ft. (*J. M. Macoun.*) Kicking Horse Lake, Rocky Mountains. (*John Macoun.*)

TARAXACUM OVINUM, Greene, Pittonia, vol. iv., p. 229.

On Sheep Mountain, Waterton Lake, lat.  $49^{\circ} 05'$ , Rocky Mountains. Herb. No. 11,711. (*John Macoun.*)

TARAXACUM LACERUM, Greene, Pittonia, vol iv, p. 230.

Canyon of the Upper Liard River, Yukon, lat.  $60^{\circ} 26'$ . June, 1887. Herb. No. 15,119. (*John Macoun.*)

TARAXACUM DUMETORUM, Greene, Pittonia, vol iv, p. 230.

A common species from Assiniboia westward to British Columbia.

TARAXACUM ERYTHROSPERMUM, Andr.

The red-seeded dandelion is probably common throughout eastern Canada, but has been seldom separated from *Taraxacum Taraxacum*. Our specimens are from Ottawa, Niagara Falls and Hamilton, Ont.

J. M. M.