

centre ; (lateral aspect), genal sulcus absent, cheek slightly shorter than the eyes, full ; eyes oval, very slightly narrowed ventrally ; (dorsal aspect), head twice as wide as long, wider than the thorax, slightly narrowed behind the eyes ; occiput immarginate, strongly concave ; ocelli arranged on an obtuse angled triangle, the distance of the lateral ocelli from the eye being about equal to that of the lateral ocelli from each other.

Thorax : Pronotum visible from above, not so wide as the mesonotum, slightly produced into a neck forward, length about half that of the mesonotum. Mesothorax slightly more coarsely reticulated than the remainder of the thorax ; parapsidal furrow distinct for half the length of the mesonotum, anterior margin rounded ; axillæ produced forward to a line drawn from tegula to tegula not meeting by half their width ; scutellum rounded, full, slightly more than half the length of the thorax ; post-scutellum narrow, strongly margined. Metathorax shorter than the scutellum, angular posteriorly ; spiracles broadly oval ; sulci absent ; lateral carinæ well marked ; median carina present ; articulation with abdomen high margined ; a distinct circular depression on either side and near to the articulation.

Abdomen : Broadly ovate, as long as thorax, greatest width at segments 3-4, subsessile ; ovipositor slightly exerted ; first visible segment the longest, about one and a half times as long as the second, the five following subequal.

Wings : Hyaline, submarginal vein slightly longer than one-third of the wing ; marginal vein the same thickness as the submarginal, as long as one-quarter of the wing ; postmarginal and stigmal veins short, subequal, the stigmal vein ending in a short knob, the knob having a short process directed towards the tip of the postmarginal vein.

Antennæ : Pubescent throughout, 13-jointed including the two ring-joints, the scape reaching almost to the median ocellus, moderately slender ; the pedicel one and a half times as long as any of the flagellar joints, obconic ; flagellar joints longer than thick, subequal, slightly and regularly clavate.

Mandibles : Brownish yellow, similar, each quadridentate.

*Male*.—Average length, 1.6 mm. Colour similar to female, reflections brighter green mixed with purple. Penis exerted. Antennæ shorter, more compact, joints of flagellum shorter than broad and more pubescent.