

558 OPERATIONS ON THE UTERINE APPENDAGES.

involved. There remains, however, a fair percentage of cases in which the disease is limited, and in which the removal of the pathological process is possible without the destruction or ablation of the entire organ.

Whatever the attitude of mind regarding these matters may be, however, the ultimate results, whether the radical or the conservative operation be undertaken, are alone of value in determining the worth of either procedure.

In order to ascertain my own experience along conservative lines, I have taken one hundred cases from my note-books, the facts and figures from which I herewith submit.

Of the one hundred women there were:—

Single	30
Married	64
Not noted	6
	<hr/>
	100

The following operations were done:—

ON THE OVARIES.

Resection.

Both ovaries in	17 cases.
Right ovary in	26 cases.
Left ovary in	13 cases.

56

Puncture.

Both ovaries in	22 cases.
Right ovary in	8 cases.
Left ovary in	14 cases.

44

In 19 women the ovary of one side was found to be so largely involved as to require removal; the right in 6 instances, the left in 13.

ON THE TUBES.

Resection.

Both tubes in	13 cases.
Right tube in	4 cases.
Left tube in	7 cases.

24

It was found necessary to remove 1 tube in 10 of the patients: the right in 4 and the left in 6 cases. The opposite side was left untouched or resected.

Results.—In the one hundred women the immediate results