

The Canadian Entomologist.

VOL. XI.

LONDON, ONT., MAY, 1879.

No. 5

DESCRIPTIONS OF NEW SPECIES OF NORTH AMERICAN BUTTERFLIES; ALSO, NOTES UPON CERTAIN SPECIES.

BY W. H. EDWARDS, COALBURGH, W. VA.

ARGYNNIS HIPPOLYTA.

Male.—Expands 2 inches.

Upper side fulvous, obscured by brown at bases of wings; the discal area of each wing lighter than elsewhere; the black markings rather heavy; the marginal lines more or less confluent; the silver spots of second row indicated on upper side by oblong spots of a pale color.

Under side of primaries pale orange-fulvous at base, and in the P-shaped spot of cell; also along the branches of median; rest of wing pale buff, except hind margin and apical area, which are ferruginous; on the sub-apical patch two silver spots, and the four or five uppermost sub-marginal spots are silvered.

Secondaries deep ferruginous, very little mottled with buff; hind margin same hue as the disk; the belt narrow, buff, much dusted with ferruginous; all the spots well silvered; those of outer row small narrow crescents, with heavy ferruginous edging to upper side; the spots of 2nd and 3rd rows small, each edged on upper side by a few scales of black; a round spot in black ring in cell, an oval in ring below cell; shoulder and inner margin silvered.

Female.—Expands 2.25 inch.

More obscured at base, otherwise like male; the basal area of primaries beneath red-fulvous; secondaries as in male, but the belt is almost lost in ferruginous.

From 3 ♂ 1 ♀ received from Mr. G. M. Dodge, and taken in Oregon, but in what exact locality is not known. Another male was received from Mr. Henry Edwards, from Northern California.