the female is more strigated and there is a tendency for the shading to form a more distinct extradiscal band than in the male. On the hind wing, however, the clouding never forms a band to the margin of the wing.

This species bears a slight resemblance to *snoviata* Packard and *puertata* Grossbeck, but differs from both in the straight extradiscal line on both fore and hind wings and in the marginal band beneath.

I take pleasure in naming this species after Mr. George H. Field, who has done so much to advance our knowledge of California Lepidoptera.

Holotype— $\sigma$ , La Puerta Valley, Calif., July 11; in my collection at the Museum of Comparative Zoology, Cambridge, Mass.; received through the kindness of Mr. Field.

Allotype—♀, La Puerta Valley, Calif., July 11, in Mr. Field's collection at San Diego. Calif.

Paratypes— $\sigma$ 's and  $\circ$ 's, from same locality, in Mr. Field's collection and my own.

## Macaria grossbecki, n. sp.

Male—Expanse 22-23 mm. Head, antennæ and legs fawncoloured. Thorax and abdomen brownish ashen, darker than in
fieldi, with a suffused appearance, without conspicuous lines or
markings as in the other species. Intradiscal line faint, pale orange,
shaded with black; mesial space heavily strigated, as in the entire
wing; not paler, as in fieldi. Discal ringlet clear black, with paler
centre. Extradiscal line faint, blackish, with slight clouding.
Veins with faint minute dots, fringe fuscous. Hind wings of the
same colour as fore wings, with a trace of an irregular extradiscal
ochreous line, blackish-shaded. Beneath paler than above, the
discal ringlets showing on all the wings. No markings apparent,
except the extradiscal brown marginal band, which shades to the
outer margin; in this respect differing from fieldi.

Female—Similar above to the male; beneath much lighter basally, where it is strigated, a brownish band running extradiscally solidly to the outer margin.

Viewed from beneath, this species has somewhat the appearance of puertata Grossbeck; but above, it appears very different