

209. *Dendroica pensylvanica*, Chestnut-sided Warbler. A moderately common migrant and breeder. It is found in deciduous second growth, and such spots in evergreen woods. In Dow's Swamp they breed yearly. Extreme dates are: May 6th to October 1st. The highwater mark in their migration is reached May 17th.

210. *Dendroica castanea*, Bay-breasted Warbler. An abundant migrant. It passes through from May 16th to June 7th (1907) and again August 28th to September 26th. This and the next species, and some of the Flycatchers, are our last migrants in spring. Its song is scarcely heard here; it is almost identical with that of the Redstart.

211. *Dendroica striata*, Blackpoll Warbler. An abundant migrant. Its chief travelling companion is the Bay-breasted. It passes through from May 17th to June 12th (1907), and again August 28th to September 26th. Song, a high, dry *tsit tsit*, repeated six to ten times.

212. *Dendroica fusca*, Blackburnian Warbler. A very common migrant and moderately common breeder. In Beaver Meadow, Dow's Swamp, Mer Bleue, Chelsea, etc., they may be seen all summer. In spring they come from the 4th to the 25th of May, reaching their climax on about the 17th; and in fall the last are seen September 9th.

213. *Dendroica virens*, Black-throated Green Warbler. A very common migrant and moderately common breeder. The habitat of this species is in mixed woods, where some hemlock occurs. Its song, *dee dee deeah dee*, announces its presence long before one sees the bird. They begin to come May 1st, attaining their highest numbers about the 17th. In fact they are most common September 10th to 27th. October 2nd marks the latest date. The Beaver Meadow waterfront is the place near Ottawa where one may look for it with the certainty of finding it in summer.

214. *Dendroica vigorsii*, Pine Warbler. A moderately common migrant and rather rare breeder. True to its name it is found in evergreens only, and nine times out of ten in pines. It breeds in the stand of somewhat taller pines along the river-shore of Blueberry Point, and in similar locations, but owing to its habitat, is rarely seen. It is found here from May 7th on.

215. *Dendroica palmarum*, Palm Warbler. The status of this Warbler will have to be revised from that given in the last list. It is a moderately common migrant only. I have taken typical examples of it only on May 10th, 1905, at Blueberry Point. Avlmer, and September 20th, 1905, near Hurdman's Bridge. The breeding birds found in our district and reported at various times in THE OTTAWA NATURALIST must go under the following