

much more dense than in *ovata*: last ventral of male small, its hind border in both sexes finely pectinate. Length 4.5–5.2 mm.

Nine specimens beaten from the flowers of the farkle-berry, *Vaccinium arboreum* Marsh, near Sanford and Ormond, March 29th.—April 6th. Easily told at a glance from *ovata* by the much more dense and coppery vestiture. Mr. Blanchard wrote me that he had had it separated but not named in his collection for more than 40 years.

*Hallomenus fuscotuturalis*, sp. nov.

Elongate-oblong. Dull brownish yellow, sparsely clothed with fine prostrate yellowish hairs; elytra with a common sutural fuscipiceous stripe which is widest in the region of the scutellum, the margins also often darker than the disk. Head finely and evenly punctate; eyes small, deeply emarginate on the inner side; antennæ with second joint one-half the length of third, joints 3–11 subequal and one-half longer than wide instead of subquadrate as in the other species of the genus except *serricornis*. Thorax at base one-half wider than long, sides gradually rounded to apex which is one-third narrower than base; disk finely and densely punctate, the basal impressions feeble. Elytra as wide at base as thorax, sides parallel for three-fifths their length, thence gradually converging to the rounded apex; their surface, as well as that of abdomen, very finely and much less closely punctate than thorax. Length 3 mm.

Six specimens beaten from scrub-oak and willow near Sanford. March 28–29, 1911.

---

NEWFOUNDLAND LEPIDOPTERA.—In a little box of insects collected at St. Anthony's during the summer of 1910, were specimens of *Argynnis freija* Thunb.; *A. myrina* Cran.; *Colias pelidne* Bdv.; *Coenonympha inornata* Edw.; *Apantesis virguncula* Kirby; *Aplectoides livalis* Smith; *Anarta cocklei* Dyar; *Mamestra sutrina* Grote; *Autographa alias* Oltol.; *Epirrita dilutata* D. & S.; *Epelis truncalaria* Walk.; *Pyrausta inaequalis* Guen.; *Crambus unistriatellus* Pack.

A. F. WINN, Westmount.

---

*Lycaena comyntas* Godt.—While collecting Geometridæ after sundown at Valcour, N. Y., July 25th., I found a male *L. comyntas* asleep on a blade of grass. Like many other "Blues," it rests for the night head downwards, the tails of the hind-wings and the black spot strongly resembling a pair of antennæ and an eye at the wrong end. A second specimen was found in the same attitude after a few minutes' search.

A. F. WINN, Westmount.