

- Myoma, necrotic, 153  
     with cardiac hypertrophy, 423  
     with hyaline degeneration of heart muscle, 423  
     with pyonephrosis, 423
- Myoma, number of, 3  
     removed by abdominal myomectomy, 517  
     obliteration of cervix with, 441  
     of umbilical depression with, 461  
     of broad ligament, 51, 53, 354, 510  
         abdominal myomectomy for, 510  
     of Fallopian tubes, 341  
     of round ligaments of uterus, 363  
     of utero-ovarian ligaments, 358  
     omental carcinoma and, 499  
     origin of, 433  
     ovarian cysts and, 344, 495  
     pain with, 448, 449  
     painful defecation associated with, 446  
         urination with, 376  
     panhysterectomy for, 593, 616  
     papilocystoma of ovaries and adenocarcinoma of uterus with, 408  
     of ovary with, 346
- Myoma, parasitic, 13. See also *Parasitic myoma*.  
     parovarian cysts with, 333  
     pear-shaped, 4  
     pedunculated, 1, 2  
     pelvic abscess with, 635, 636  
         molds and, 6  
         peritonitis with abscess formation and, 344  
     perirectal abscess with, 391  
     phleboliths in vaginal wall with, 440  
     polypi of cervical mucosa with, 298  
         of uterus associated with, 325, 328, 329  
     pregnancy and, 467, 473, 477, 507, 643  
     pressure of, on bladder, 365  
         on iliac veins, 234  
         on ureters, 381  
     pains with, 448  
     prolapse of rectum with, 391  
         of uterus with, 460  
     pruritus ani associated with, 446  
         vulva with, 444  
     pyelonephrosis with, 397  
     pyometra associated with, 310  
     pyosalpinx with, 338, 340, 396  
     rectal diverticula, rupture of, and, 500  
     rectum and, 386  
     removal of, in apparently inoperable cases, 636  
     renal colic with, 466  
     resemblance of, to fetus, 6  
     respiration and, 450  
     results of operations for, 671
- Myoma, retention cysts in kidney with, 427  
     of urine with, 376  
     retraction of umbilicus with, 461  
     retroperitoneal, development of, 631  
         sarcoma and, 501  
     sacculization of bladder with, 370  
     sacral markings with, 5  
     saddle-bag appearance of uterus with, 5  
     salpingitis with, 338  
     sarcoma developing from, 169, 172, 173, 180, 208, 420  
         of bladder associated with, 420  
         of uterus and, 492  
     secondary carcinoma of uterus with, 295
- Myoma, sessile, 1, 2  
     shape of, 4  
     stimulated by chorio-epithelioma, 482  
         by extra-uterine pregnancy, 477  
         by hydatidiform mole, 480  
         by omental metastases, 499  
         by retained menstrual flow, 485  
         by retroperitoneal sarcoma, 501  
         by simple ovarian cysts, 497  
     size of, 3  
     small developing in uterine mucosa, 333  
         unattached to uterus and situated between tube and ovary, 364  
     snugly filling pelvis, rendering abdominal hysteromyomectomy difficult, 631  
     sterility associated with, 457
- Myoma, submucous, 1, 59, 571. See also *Submucous myoma*.
- Myoma, subperitoneal, 1, 45, 81, 134, 249, 444, 508
- Myoma, subvesical, 365, 510
- Myoma, suppurating, 134-153  
     cause of, 152  
     interstitial, 144  
     intraligamentary, 137  
     opening into cecum, 45  
         into colon, 142  
         into small bowel, 47  
     subperitoneal, 134, 135, 139  
         communicating with bowel, 134  
     suspicion of carcinoma of cervix with, 302, 304  
     symptoms of, 434-459  
     temperature with, 457  
     thickening of broad ligaments with, 355  
         of left lobe of liver with, 464  
     three-leaf-clover-shaped, 5  
     thrill felt on vaginal examination with, 442  
     thrombosis of veins in mucosa with, 320  
     torsion of cervix with, 77  
         of uterus with, 77, 79  
     transition of, into sarcoma, 182