ARCTOMYS MONAX, Linn.

0

¢.

x.

of

з.

ıt-

Woodchuck. Ground-hog. Arctomys empetra, (Sabine) Rich. Arctomys pruinosus, Gmelin. A-kwa-kwa-chis. Saulteaux.

Nova Scotia and New Brunswick and westward around the shores of Hudson's Bay to the Mackenzie and Liard Rivers, as far north as Lat. 62°.

ARCTOMYS CALIGATUS, Eschscholtz.

Hoary Marmot.

Arctomys pruinosus, (Gmel.) Rich. Arctomys Okanaganus, King.

Rocky Mountains westward to the Cascade Range and northward to the Arctic Circle. Often seen among heaps of angular masses of loose rock. I can find no reference to its being found east of the Rocky Mountains.

The specific name caligatus has here been used instead of pruinosus, which latter name, as was pointed out to me by Prof. Macoun, appears to have been in all probability first applied to a light colored specimen of monax. Pennant's description on which Gmelin's name pruinosus was founded, as given in Coues and Allen's "Monographs of North American Rodentia" p. 926, agrees so well with a specimen of A. monax from Hudson's Bay, the original locality, now in the Geological and Natural History Museum, that I have very little hesitation in placing Gmelin's species as a synonym of A. monax; and although the A. pruinosus of Richardson is undoubtedly the true A. caligatus, his name cannot stand on account of its having been formerly applied to a different species.

## INSECTIVORA.

SCALOPS AQUATICUS, Linn. Common Mole. Common throughout Eastern Canada.

2o