

12, on whom he had operated, and restored the anterior wall by Maury's operation. A large oval flap was first taken from the perineum and fixed beneath a short flap dissected from above. After union had taken place, the sides of the flap, which were unattached, were further dissected down and fixed beneath short lateral flaps. In the first operation, a hole had been made in the centre of the perineal flap for the rudimentary penis. The parts all united well except at the upper part, where a small portion sloughed and allowed urine to exude, and so prevented skin-grafting being to any large extent successful. This fistulous opening had, however, been closed by a recent operation, and now the bladder was completely covered and the parts had all skinned over. The boy was able to retain a couple of ounces of urine, and the double hernia which had previously existed as the parts contracted was completely cured.

DR. RODDICK considered the operation admirable. He had operated on a young woman some years ago for extrophy of the bladder, and had selected Ayer's method. A large square flap had been dissected from the abdomen above the bladder and turned down with the cuticular surface innermost. The raw surface was subsequently covered over by lateral flaps. The operation thus far had proved very successful. The patient left the hospital with the intention of returning in a few weeks to have the operation completed. She failed to do so. It was learned that she had got married!

*Study of Koch's Treatment in Berlin.*—DR. G. T. ROSS read a paper on the above subject, which appears in the present issue of this JOURNAL.

Brief reports of cases submitted to Koch's treatment in the Montreal General Hospital, which are being published in this JOURNAL, were made by Drs. Roddick, MacDonnell, Jas. Bell and Johnston.

DR. RODDICK was not yet convinced that the results would be as good as predicted. He agreed with Dr. G. T. Ross that the remedy was a dangerous one, and that all experiments should be made in hospital.

DR. R. L. MACDONNELL stated that, recognizing the respon-