

382. *LOLIUM*, L. - - DARNEL. RYE-GRASS.
 883 *L. PERENNE*, L. Near railway depot, Carleton, *Hay*. No specimen.
 884 *L. TEMULENTUM*, L. Near railway depot, Carleton, *Hay*. Almost the only noxious grass known. No specimen.
383. *TRITICUM*, L. - - - - WHEAT.
 885 *T. repens*, L. A very troublesome weed on sandy soil. Varies greatly, especially on the sea coast.
 886 *T. caninum*, L. On intervals, Coal Branch, Kent Co.
384. *HORDEUM*, L. - BARLEY. SQUIRREL-TAIL GRASS.
 887 *H. jubatum*, L. Sandy soil along the Gulf shore.
385. *ELYMUS*, L. - LYME-GRASS. WILD-RYE.
 888 *E. Virginicus*, L. On intervals, Hampton, Coal Branch.
 889 *E. Canadensis*, L. University Herb. Sussex and Petiteodiac,
Brittain; E. River, Carleton Co., *Hay*.
 890 *E. mollis*, Trin. Sand beaches, Kouchibouguac, *Fowler*; Restigouche, *Chalmers*.
386. *ASPRELLA*, Willd. - BOTTLE-BRUSH GRASS.
 891 *A. Hystric*, Willd. (*Gymnostrichum Hystric*, Schreb.) Eel River, Carleton Co., *Hay*.

CRYPTOGAMIA.

CLASS III.—ACROGENOUS CRYPTOGAMS.

XC. EQUISETACEÆ.

387. *EQUISETUM*, L. - HORSETAIL. SCOURING RUSH.
 892 *E. arvense*, L. Very common in damp soil.
 893 *E. pratense*, Ehrh. Restigouche River, *Chalmers*.
 894 *E. sylvaticum*, L. Very common throughout.
 895 *E. limosum*, L. Rather common. Kent, *Fowler*; Tobique Lakes, Carleton, *Hay*, Herb.; Nepisiquit Lakes, *Bailey*, Can. Nat. 1864, p. 90.

- 896 *E. hyema* Viet
 897 *E. scirpo* Le N
 388. POI
 898 *P. vulgare* the n
 River
 389. ADI
 899 *A. pedaturn*
 Kesw
 II. 3
 Daley
 390 PTEI
 900 *P. aquilina*,
 391. PELI
 901 *P. gracilis*,
 Morris
 Wood
 392. ASPI
 902 *A. viride*, H
 ter Co
 Fowler
 Herb.
 903 *A. thelypter*
 Fowler
 toria C
 904 *A. Filix-fœm
 393. SCOL
 905 *S. vulgare*,
 Herb.
 fax, Bu
 394. PHEG
 906 *P. polypodioides*
 Herb.*