

Hind wing paler gray than fore wing, slightly darker around termen, cilia paler, with a faintly darker basal line. Under side the same, a black dot on costa before apex. Under side fore wing darker gray, with a few black dots on costa before apex.

Four specimens, Rounthwaite, Man., July 11 and 12, L. E. Marmont.

Co-type in U. S. Nat. Mus., Cat. No. 9870. I am indebted to Mr. Busck for correcting my generic determination of this species.

*Mompha Claudiella*, sp. nov.—Expanse, 20 mm.

Head and patagia white, with a faint rosy tinge; palpi white, with a few dark scales on outside of tuft of second joint; antenna fuscous, with a whitish bloom; thorax leaden-black in centre, narrowly edged in front with rosy-white; on posterior half there is a patch of deep rosy-white, and the middle scales at their posterior end are of the same colour; abdomen gray, anal tuft with an ochreous tinge; legs cream-white, heavily shaded with black in front.

Fore wing white, with a faint rosy tinge, shaded with bluish-gray on inner fifth of costa, and a shade of the same colour through middle of wing from inner tuft to end of cell. A small shade of this colour on outer third of costa, and a similar one below it on dorsal margin. The upper half of termen and apex are shaded with gray, in which are sprinkled whitish, black and brown scales; there is a dark brown dot in cilia at apex, and two before apex in costal cilia, and one below apex in the long cilia; the outer ends of the apical cilia are gray, otherwise the cilia is ochreous. The tufts are very large and high; the three most prominent ones are white towards base and ferruginous at their outer ends. There are two close to base of wing, one in middle of wing, and one below it and obliquely outward from it close to dorsal margin; before the latter, and in a streak following the former, are broad black scales; the third is between fold and dorsal margin in middle of wing; it is also followed by a patch of broad black scales from its upper end. There is a smaller but similar tuft at outer fourth of wing, on lower margin, and a small tuft above the third large one on upper half of middle of wing. Below costa, at inner third, is a small tuft of black scales.

Hind wing dark gray, cilia ochreous, the same shade as cilia of fore wing. Under side both wings smoky black, with a whitish streak through middle of hind wing, and the apex of both wings whitish.

Three specimens, Rounthwaite, Manitoba, July 11 and 12, collected by Claud Marmont, whose name I am glad to give to this very beautiful species.

Co-type in U. S. Nat. Mus., Cat. No. 9871.