

***Pandion haliaetus carolinensis* (Gmel.) OSPREY.**

Rare, except in May and June, when they are frequently observed along the Gaspereau River following the movement of the ascending fish.

***Asio wilsonianus* (Less.) LONG-EARED OWL.**

Common except in winter.

***Asio accipiterinus* (Pall.) SHORT-EARED OWL.**

Common on the Grand Pré from April till the middle of December.

***Syrnium nebulosum* (Forst.) BARRED OWL.**

Fairly common in heavily wooded regions.

***Nyctala acadica* (Gmel.) SAW-WHET OWL.**

Present throughout the year, but most common in winter.

A nest was taken last April containing six eggs.

***Bubo virginianus* (Gmel.) GREAT HORNE-OWL.**

Uncommon. Found in densely wooded districts.

***Nyctea nyctea* (Linn.) SNOWY OWL.**

Rare winter visitant.

***Surnia ulula caparoch* (Mill.) HAWK OWL.**

Rare. An example was taken last October.

***Coccyzus erythrophthalmus* (Wils.) BLACK BILLED CUCKOO.**

Fairly common from May 20th till September.

***Ceryle alcyon* (Linn.) BELTED KINGFISHER.**

Fairly common about suitable places from May till October.

***Dryobates villosus* (Linn.) HAIRY WOODPECKER.**

Fairly common throughout the year.

***Dryobates pubescens* (Linn.) DOWNY WOODPECKER.**

Common throughout the year.

***Picoides arcticus* (Swains.) ARCTIC THREE-TOED WOODPECKER.**

Rare winter visitant.

***Sphyrapicus varius* (Linn.) YELLOW-BELLIED SAPSUCKER.**

Fairly common summer resident.

***Cephalæus pileatus* (Linn.) PILEATED WOODPECKER.**

Rare. One was taken in September, 1897.