

City	Expend.	Estimated Population 1927	(2) General Death Tuberculosis	Infant Mortality Rate	Total Deaths Diarrhoea & Enteritis 0 - 2 years	
	Per Capita 1927		Rate, (Deaths per 1,000 Population) 1927	Deaths, 0-1 year per 1,000 live births 1927		
1. New York	.72	5,970,000	11.8	86	56	729
2. Chicago	.74	3,101,000	11.5	82	62	547
3. Philadelphia	.50	2,036,000	12.1	85	64	290
4. Detroit	1.01	1,341,385	10.8	91	70	296
5. Cleveland	1.01	984,000	9.6	80	56	115
6. St. Louis	.65	830,000	12.9	62	56	123
7. Baltimore	.79	813,333	14.1	87	81	191
8. Boston	1.06	793,000	14.0	86	76	230
9. Pittsburg	1.18	637,000	13.4	73	71	114
10. Buffalo	.92	542,060		figures not yet available		
11. Newark	1.02	467,000	10.9	(1) 82	63	78
12. Rochester	.96	317,000	11.1	(1) 67	63	43
MONTREAL	.39	699,500	14.9	126	113	870

(1) Includes deaths of citizens in sanatoria outside city

(2) Includes all deaths occurring in city; non-resident deaths not excluded.

These rates were obtained from the cities by correspondence.