

1692. *Leptinus testaceus*, Müll.—From a living shrew, September 9, 1892.
- 1701a. *Necrophorus Melsheimeri*, Kirby.—Abundant from May onwards.
1702. *Necrophorus vespilloides*, Hbst.—Occasional.
1706. *Silpha lapponica*, Hbst.
1715. *Necrophilus hydrophiloides*, Mann.—Abundant, especially during winter. Active under carrion even during snow.
1716. *Pelates latus*, Mann.—Abundant under logs and loose bark. Seen *in cop.* February 16th.
1719. *Agyrtes longulus*, Lec.—January 3. In flood refuse. Several taken.
1720. *Sphaerites glabratus*, Fab.—Common during summer, under carrion.
1722. *Pinodytes cryptophagoides*, Mann. From June to September, under logs.
1724. *Catoptrichus Frankenhauseri*, Mann.—Somewhat rare. Under carrion in autumn.
1725. *Choleva egena*, Horn.—Two skimmed from surface of flood refuse in meadow, 14th Oct., 1893.
1726. *Choleva luridipennis*, Mann. (also one specimen of a black variety). Abundant during autumn, under carrion.
1741. *Ptomaphagus pusio*, Lec.—Rare in hen's nest, 6th November, '91. *Hydnobius*, probably n. sp.—In rotten grass, 17th June, 1892.
1769. *Anisotoma humeralis*, Horn.—January 3. In flood refuse. A few taken.
1803. *Agathidium concinnum*, Mann.—Under bark in December. Not common.
1804. *Agathidium rotundulum*, Mann.—Rare. Under carrion. January 3. In flood refuse. Abundant.
1807. *Agathidium pulchrum*, Lec.—Rare. Taken from mud gallery made by ants on trunk of dead alder.
1847. *Scydmaenus biformis*, Makl.—Not common. In moss at tree roots in February.
1899. *Batrissus frontalis*, Horn.—Not common. In ants' nests.
1911. *Bryaxis albionica*, Mots.—Occasionally in winter, in moss at tree roots.
1955. *Faronus parviceps*, Makl. — One specimen taken on wing in September.
2025. *Homalota granularis*, Mann.—Common in sea-weed.