

## LIST OF FIGURES (continued)

<u>Figure Number</u>		<u>Page Number</u>
3-14	Geographic distribution of pH levels measured in Adirondack Lakes higher than 610 meters elevation, June 24-27, 1975.	3-55
3-15	Comparison of lake pH, 1930 to 1975, for the Adirondack Mountains area of New York.	3-56
3-16	Cumulative comparison of historic and recent pH values for a set of 138 Adirondack Lakes, New York.	3-57
3-17	Surface water alkalinity of New England states.	3-59
3-18	New Jersey stream pH, 1968-1979, Oyster Creek and McDonalds Branch.	3-60
3-19	Distribution of pH of North Carolina streams June, July, August 1960-1963 (upper), compared with June, July 1979.	3-62
3-20	Distribution of alkalinity of North Carolina streams June, July, August 1960-1963 (upper), compared with June, July 1979.	3-63
3-21	Snowmelt-runoff hydrogram and changing water quality for Filson Creek, Minnesota, Boundary Waters Canoe Area Wilderness region.	3-65
3-22	Discharge, hydrogen ion load per unit area, pH, and depth of precipitation for each day that a precipitation event occurred for Harp Lake No. 4.	3-67
3-23	Hydrogen ion content of streams draining Red Chalk Lake watersheds No. 3 and No. 4 (Dorset, Ontario) showing effects of a 2 cm rainfall (pH 4.06) between 11:00 p.m. July 12, 1977 and 3:00 a.m. July 13, 1977.	3-68
3-24	Mean daily pH for the Shavers Fork River at Bemis W. Virginia and precipitation event pH and accumulation at Arborvale, W. Virginia.	3-70
3-25	Relative number of taxa of the major taxonomic groups as a function of pH.	3-72