heard from him with pleasure that he is going to give a number of varietal forms in the new volume now being published; the plates will be for the most part figured from examples captured by me in 1883-4.)

C. Scudderii, Reak.

2 Slow of movement, like Christina, and easily captured.

C. Occidentalis, Scud, including southern form Chrysomelas, Hy. Edw.

A lively insect, only taken when the sun was hottest and the day brightest.

C. Edwardsii, Behr.

No notes taken.

C. Alexandra, Edw.

2 very lively, short flights; 2 not observed.

C. Eurytheme, Boisd.

Winter form Ariadne, Edw.

" Keywaydin, Edw.

All the different forms of this insect are the liveliest I came across in the North West Territories.

C. Hagenii, Edw.

Very like the above, and almost impossible to catch on the wing, from the zig-zag movement in flying, and its long flights, often flying high in the air. Was taken in 1886 by me, at Kakabeka Falls, Lake Superior.

- C. Philodice, Godt.
- C. Interior, Scud.

Apparently a lazy insect; easy of capture. Very few taken in 1883 and 1884.

C. Elis, Streck.

A has a remarkably short flight, dropping suddenly to the ground, getting itself entangled amongst the grass and foliage; not alighting, as most Coliads do, where they can be seen, on a leaf or on a flower.

C. Elis is a new species described by Mr. Herman Strecker, after my return from the Rocky Mountains, in 1884. At present nothing is definitely known about the male of this species, although two forms of the