

Scutellum black, with the tip, the lateral margins anteriorly, and a median line reaching only to the transverse impressed line, yellow. Elytra white, dusky toward the apex, marked as in *variata*, with oblique brown bands forming a lozenge anteriorly, including a pale commissural spot, and an x posteriorly with the tip of the clavus as its centre, the median bands being common to both marks. These brown bands do not attain the costal margin, and are more strongly distinguished than in *variata*. Abdomen and pleural pieces deep black, the margin of the tergal pieces slenderly yellow, venter sometimes yellow with the segments and connexivum bordered within their margins with blackish. Legs brown, the joints and tibial spines pale.

New York; New Hampshire. Described from three examples; one male taken by me at Lancaster, N. Y., July 12th, 1889, a female taken at "High Bridge," New York City, in June, by Mr. E. B. Southwick, and a second female taken on the summit of Mt. Washington by Mrs. Annie Trumbull Slosson, to whom I take pleasure in dedicating this pretty little species as a slight tribute to her persevering devotion to the study of entomology.

### 3. CHLOROTETIX NECOPINA, *n. sp.*

Form and size of *tergatus* nearly. Vertex with a black transverse band between the eyes. Colour above greenish-testaceous, marked and clouded with brown; elytra fuscous, with strong white nervures. Length, 7 mm.

Vertex a little more produced than in *tergatus*, and the front proportionately longer and narrower; ocelli contiguous to the eyes. Last ventral segment of the female thin, broadly and deeply cleft nearly to its base; this incisure at its apex armed with a small blunt tooth, the lateral lobes ovate, in one example feebly angled at apex. Pygofers stout, very slightly exceeded by the oviduct, and armed with a few stout brown bristles. Other characters about as in *tergatus*.

Colour: Beneath pale greenish tinged with testaceous, especially on the front, which is marked on the sides by about ten nearly obsolete brown arcs, and two brown dots are at the base of the clypeus; eyes dark brown edged with pale. Vertex with a broad transverse blackish band just behind the ocelli. Pronotum testaceous, pale before and on the median line, either side of which is a brown cloud, and a smaller one occupies the latero-posterior angle, transverse wrinkles less distinct than in *tergatus*. Scutellum marked with a piceous triangular spot within the