173. Leptura behrensii LeConte.

Rare Two specimens taken by Baumberger in Northern Sonoma Co.; type specimen from Mendocino Co.; one taken by Van Dyke in Humbolt County.

174. Leptura scripta LeConte.

Common on Coast belt from Marin County to northern line on flowers, especially Azalea, Rhododendron; rare in the various parts of the Sierras. June 9.

175. Leptura gnathoides LeConte.

Found in Sierras and Northern California. Rare. Taken in Placer County by Van Dyke on young trees of post cedar.

176. Leptura valida LeConte.

Found rarely from Shasta to the Southern Sierras. Breeds in pine. Taken by author at Donner Lake. July 18.

177. Leptura insignis Fall.

Taken by Fuchs and Fenyes at Monterey. Breeds in Monterey pine; remains also found in *Pinus muricata* at Inverness; a specimen was bred by Van Dyke from Douglas spruce branch of unknown locality.

178. Leptura aspera LeConte.

Northern, may possibly run over into California.

179. Leptura cubitalis LeConte.

This species, which looks like an Acmæops, is fairly common in the middle part of California. It is found on wild rose, etc., in Sonoma, Napa, San Mateo, and Lake Counties. June 8–29.

180. Ophistomis ventralis Horn.

Taken at El Taste, Lower California.

181. Strangularia delicata LeConte.

Occasionally found in most parts of the State, but is more common in the north and middle Sierras on various flowers. June 3.

(To be continued)

Mailed July 10th, 1918.