solution at each injection. At the same time, two or three times a week, a tampon impregnated with equal parts of alum, tannin and salol should be introduced, and every evening a vaginal suppository containing four grammes (3j) of tannin or iodoform, one-half a gramme (7½ grs.) of glycerine and cacao butter.

The writer concludes, after two operations of Symphyseotomy and some investigations on the cadaver, that not more than two centimetres separation of the pubic joint follows simple division of the articulation, while by section of the ligamentum arcuatum inferius the joint will separate to the extent of 5 to 6 ctm.—

(DODERLEIN, in Med. Record.)

— A mixture of chloroform (ten parts), ether (fifteen parts) and menthol (one part), used as a spray, is recommended as an excellent and prompt means for obtaining LOCAL ANESTHESIA lasting about five minutes.—(Boston Med. and Surg. Journal.)

-Hayem* gives the following prescription

for the relief of Acute Coryza:

R. Acid. carbolic.,

Aquæ ammoniæ, āā z ijss Alcohol, z v Aquæ destil, z j. M.

*Rev. de Laryngol., d'Otol. et de Rhinol., in Boston Med. and Surg. Journal.

Sig.—Inhale from several drops upon a piece of bibulous paper.

—Lotion for Pruritus Vulvæ (Practi-

tioner)

B. Hydrargyri perchlorid., gr. j.
Aluminis, gr. xx
Pulv. amyli, 3 jss
Aquæ menthæ pip. q. s. ad 5 vj.

M. et fiat lotio.

Sig.—Apply externally to the affected parts. -Dr. Carasso Michele, Director of the Military Hospital at Genoa, has used since 1888, in the treatment of pulmonary tuberculosis, constant Inhalations of Oil of Peppermint (Boston Med. and Surg. Journal, Jan. 11th, 1894). He combines the inhalation with the internal administration of an alcoholic solution of creosote, glycerine and chloroform, to which is added oleum menthæ piperitæ, 1: 100. His results are reported as remarkable. only were incipient cases cured, but advanced cases also, some thirty-nine in all, with cavityformation and abundant bacilli in the sputum. All the cases treated were of pulmonary diseases only, without tubercular affection elsewhere.

EPITHELIOMA of superficial variety may be treated with alternate applications of ten per cent, methyl-blue and twenty per cent, chromic

acid.—(Darier, in Med. Record.)

—The heart cannot be weakened under Chloroform except by interference with the breathing. It is useless and dangerous to take the pulse as a guide. Wafch the respiration.

Safety is insured only by regular natural breath ing.—(LAWRIE, in *Med. Record.*)

—Indications for Cholecystotemy are frequently recurring biliary colic without jaundice, where medical treatment has failed. Persistent jaundice where the onset is ushered in with pain, and where recurring pains, with or without ague-like attacks, render it probable that the cause is gall stones in the common duct. Distended gall-bladder from impaction of calculi in the ducts. Empyema of the gall-bladder. Persistent jaundice with enlargement of the gall-bladder dependent on some obstruction in the common duct, even where the cause cannot be clearly made out.—(Robson, in Med. Record.)

—Supra-Sphincteric Ulcers of the Rectum require the galvano-cautery under anæsthesia and with the aid of speculum. Boric acid lotions and iodoform tampons may be used in the after-treatment, and antiseptics, such as naphthol, administered internally.—(Quenn in

Med. Record.)

в.

—FOR ROSACEA, Petrinio in Medical Re-

Ichthyol, 2 parts
Resorcin, 1 part
Collodii flexil, 30 parts.

—Dr. Liveing (Münchener Med. Wochenschrift, in Cinn. Lancet-Clinic, Jan. 13, 1894) recommends in Itching of the Anus from Pin Worms the following salve, to be rubbed in every evening:

B. Mercurial oint., equal parts.

Vaseline,

For the same affection the following formula is also of service:

Calomel, gms. 3
(grs. xlv).

Vaseline, gms. 30
(grs. 3 j).

Cocaine, 1 part
Bismuth subnitrate, 2 parts
Lanoline, 20 parts

—In Cancer of the Uterus total extirpation is destined to be the operation whether the disease is cervical or corporeal, and if done in time will permanently cure a certain portion of cases. If any tissues adjacent to the uterus are already involved, it should not be attempted.—(Lewis in *Med. Record*).

—Dr. Fitch (Charlotte Medical Journal) recommends the following in SUMMER COM-

PLAINT:

B. Acid. hydrochloric. dilut., mxvj
Pepsin. pur., 3s
Bismuth. subnitrat., 3 ij
Syrup., f 3 ij
Aquæ destillat., f 3 ij.

Sig.—Shake the bottle, and give a teaspoonful before each feeding or nursing to an infant one year old, half the dose to an infant six months.

This mixture must be made fresh every