

often mentioned and figured in agricultural reports (first by Glover, U. S. Agr. Rep., 1855, p. 95, pl. viii., fig. 9) as *M. femoratus*, but it is nevertheless found in just such situations as are given for the latter, and is the only species I met with in the South. The locality of this species is given in the Check List as the Western States.

Leptoglossus phyllopus Linn. Numbers taken on thistles and in much the same places as the preceding. N. O. ; B. la F.

LYGAEIDÆ.

Myodocha serripes Oliv. One taken under old wood?

Melanocoryphus bicrucis Say. Four taken on plants in May. B. la F.
This species I have also taken in Kansas and Michigan.

PYRRHOCORIDÆ.

Largus succinctus Linn. Fifteen taken in April on leaves of young shrubs of elder on a plantation. *In coitu*, 21st April. B. la F. This is no doubt the species referred to and figured by Glover (U. S. Agr. Rep., 1855, p. 94, pl. viii., fig. 7) as the "Red-edged-winged Reduvius," as his description applies well to this insect.

PHYMATIDÆ.

Phymata erosa H. Schf. Three taken on some roadside weeds in May. B. la F. The locality of this species was formerly given as Mexico.

REDUVIDÆ.

Zelus bilobus Say. One taken 22nd May on ragweed. B. la F.

Apiomerus sp. A larva taken under old wood?

Sirthenea carinata Fab. Three taken in April under logs in damp places on the edge of the swamp. N. O.

Rasahus biguttatus Say. Four nymphs taken in April under logs in same places as the preceding species. N. O. ; B. la F. The locality of this species was formerly given as the Western States.

Melanolestes picipes H. Schf. Six taken in April in same places with the two preceding. N. O. Have also taken this species in Kansas.

Conorhinus variegatus Drury. One nymph taken with the preceding species. N. O.

GALGULIDÆ.

Galgulus oculatus Fab. One taken in slow water in April. N. O.