

Primaries ochreous, shaded with darker, costa and inner margin gray; a narrow black basal dash inclined slightly upward; ordinary spots prominent, orbicular gray, filled with darker shade, and outlined with black, except basally, where the black marginal lines proceed for a short distance parallel to each other towards the base. Reniform well scalloped outwardly, margined with black opposite orbicular, the space between the two spots slightly darker than ground colour. Claviform outlined in black; a submarginal row of black dashes in interspaces, most prominent in central portion of wing; marginal area considerably darker than remainder of wing, and terminated by fine black line, slightly lunate in interspaces, second anal vein black from base to margin; cubitus gray as far as reniform; all veins blackish in submarginal area, veins  $M_3$  and Cu being prominently bordered with gray. Fringes smoky, with a fine yellow basal line and a darker median shade. Secondaries white at base, with a broad smoky-brown border and a brownish lunule at end of cell, fringes white. Beneath, primaries yellowish-white, with spot at end of cell, and costa and outer margin sprinkled with brown; secondaries white, costa with brown sprinkling; slight spot at end of cell.

Expanse, 33 mm.

Habitat.—Kerrville, Texas, 2 ♀ ♀'s.

Type, coll. Barnes.

This species is allied to *E. Hollemani* Grt., is, however, considerably lighter in general appearance; the reniform and orbicular are distinctly separate, whereas in typical *Hollemani* (we possess specimens compared with type) they are fused; the claviform is also present in our species, and the black basal dash less prominent.

*Stibadium mavina*, n. sp.

♂.—General colour ochreous, moderately frosted with white and brown scales. T. a. line scarcely perceptible, marked on costa at about one-third of the length by a slight patch of whitish scales. T. p. line narrow, white, angled outwardly near costa, thence oblique and nearly parallel to margin. Median space brighter yellow, contrasting especially with submarginal portion of wing. Ordinary spots practically indistinguishable, very faintly outlined with white. Slight whitish mark proceeding obliquely downwards from costa near apex. Secondaries lighter than primaries at base, shading into darker towards margin. Fringes and thorax concolorous with wings. Beneath, without markings, sprinkled with darker scales along costa and outer margin, especially on secondaries.

Habitat.—Provo, Ut. (Spalding).

Type, 1 ♂, coll. Barnes.