(or three hundred cubic centimetres) of water for fifteen minutes; cool, filter and wash the filter with sufficient water to make seven ounces (or three hundred and fifty cubic centimetres). In this dissolve the sugar with the aid of heat, and strain. When cold, add the solution of phosphates and sufficient water to measure twenty fluid ounces (or one thousand cubic centimetres).

Note.-Large amounts used.

SYRUPUS HYPOPHOSPILITUM COMPOSITUS.

(Compound Syrup of the Hypophosphites.)

Calcium Hypophosphite	80	grains.	12	grammes.
Potassium Hypophosphite	40		6	• "
Manganese Hypophosphite.	40	• 6	. 6	
Iron Hypophosphite	40	"	6	"
Potassium Citrate	30	**	4.5	• •
Strychnine Hydrochloride	2	"	0.3	4.
Quinine Hydrochloride	8	**	1.2	"
Sugar	14	ounces.	700	"
Water	\mathbf{A}	sufficient	quan	tity.

SYRUPUS SENEGÆ.

(Syrup of Senega.)

Liquid Extract of Senega	4 ounces.	200 ccs.	
Sugar Water	14 ''	700 grammes.	
Water	A sufficient quantity.		

Mix the Liquid Extract of Senega with ten ounces (or five hundred and fifty cubic centimetres) of water; filter through Kaolin, washing the filter with distilled water; dissolve the sugar in the filtrate; strain and add water to make twenty fluid ounces (or one thousand cubic centimetres) of the Syrup.

Dose.— $\frac{1}{2}$ to 1 fluid drachm.

Note.—Preferred to tincture.

SYRUPUS IPECACUANHÆ.

(Syrup of Ipecacuanha.)

Liquid Extract of Ipecacuanha.	1 fld. oz. 50 ccs.
Acetic Acid	96 minims, 10 "
Glycerine	2 ounces. 100 "
Sugar	14 " 700 grammes.
Water	A sufficient quantity.

Mix the Liquid Extract of 1 pecacuanha, Acetic Acid and ten ounces (or five hundred cubic centimetres) of water; filter through Kaolin into a vessel containing the Glycorine; add the sugar and dissolve without the aid of heat; strain and add water to make twenty fluid ounces (or one thousand cubic centimetres) of the Syrup.

Dose.-1 to 2 drachms.

Note.—Strongly recommended.

TINCTURA ARNICÆ FLORES.

(Tincture of Arnica Flowers.)

Moisten the powder with four fluid ounces (or two hundred cubic centi-