

diet some degree of Anemia is very likely to supervene. This is especially true after a long-continued Typhoid, as well as in cases of chronic nephritic disease, in which milk is the principal or exclusive food. The occurrence or such a "milk diet" Anemia seems to be, in many instances, responsible for a tardy and protracted convalescence. Such iron-poverty can be prevented by administering Pepto-Mangan (Gude) both during and after the milk diet period, thus supplying the essential iron in the most easily tolerable, non-irritant and promptly assimilable form. This palatable organic, ferruginous compound is entirely free from disturbing effect upon the digestion, and does not irritate, or constipate, nor does it in any way interfere with such other treatment as the physician may see fit to adopt.

### CONSTANTLY FAVORABLE RESULTS.

Dr. John Arthur Diggle, Med. Ref. Globe Accident Assur. Soc., of London, Eng., in writing of antikamnia tablets, says: "I may state at the outset that they satisfied me well and the

**DOCTOR'S**  
**BRASS SIGNS**  
 & RUBY GLASS SIGNS  
**G. BOOTH & SON**  
 21 Adelaide St. W., Toronto

## Glyco-Thymoline

IS INDICATED FOR

## CATARRHAL CONDITIONS

**Nasal, Throat, Intestinal,  
 Stomach, Rectal and  
 Utero-Vaginal**

SAMPLES ON APPLICATION

**KRESS & OWEN COMPANY**  
 210 Fulton St., & NEW YORK

