

foveæ fairly deep ; colour reddish brown with the carinæ and margins light yellowish brown.

Front about twice as long as wide, sides very slightly arcuate, widest a little below the eyes ; clypeo-frontal suture slightly curved, median carina distinct, forked just below the apex of the head ; colour uniform reddish brown except a narrow strip along the clypeo-frontal suture, which is light yellow.

Clypeus uniformly black with the median carina fairly distinct.

Second segment of the antennæ a little more than twice as long as the first segment, second segment roughened by tubercles, proximal segment black, second segment yellowish brown.

Prothorax not quite so deep as the vertex, carinæ distinct, the lateral carinæ quite widely divergent, fading out just before the posterior margin, posterior margin distinctly concave ; colour reddish brown, except the narrow posterior edge, which is almost white.

Coxæ light yellow ; femora and most of the tibiæ reddish brown, tips of the tibiæ and the first two tarsal segments very light yellow, last tarsal segment dark brown, spur almost white with large, prominent, black serrations.

Scutellum almost twice as deep as the prothorax, sides very decidedly arcuate, tip a rounded point, carinæ all distinct, lateral carinæ reaching the posterior margin of the scutellum and curving outward ; colour dark brown, except the tip, which is light yellow, almost white.

Elytra extending one-third their length beyond the tip of the abdomen, veins brown, membrane smoky ; abdomen uniformly dark brown.

Genitalia : Pygofer ending in a blunt point, reddish brown, ovipositor reddish brown at the tip, lighter at the base, plates dark within, lighter on the free edge, apex very gradually curved, extending about one-third the length of the pygofer, style light yellow. Length, including elytra, 3 mm.

Macropterous ♂.—Same form, size and general characters as the female. Genitalia : Opening of the pygofer broad oval, superior wall of anal tube prolonged into long, incurving, tusk-like horns, the points resting in indentations of the inferior wall of pygofer, styles large hook like organs with the hooks pointing outward ; colour dark reddish brown.

Described from a pair taken by the author at Middletown, N. Y., July 12, 1910. Two females of this species were taken at the same locality on July 11 and 18, 1910. Types in the author's collection.