

half concealed by the pronotum; apical limbus very broad and corrugated. Abdomen black; legs yellow.

Type.—Female.

Length 3.4 mm.; width between humeral angles 2.2 mm.

Locality.—Peru.

12. ***Tragopa decorata*** sp. nov. (Figs. 12 and 13).

About the size of the preceding. Apparently near *T. fulvovaria* Fairm., in its decorations; but smaller, not black, and without the large posterior spot of Fairmaire's species. The head shows no markings.

Decidedly longer than wide; beautiful rich chocolate brown shading to light brown on shoulders and head; decorated with numerous irregular yellow dots and bars bordered with dark brown. Head slightly wider than long; uniform light brown; front convex; ocelli yellow, farther from each other than from the eyes; eyes black bordered with yellow. Pronotum chocolate brown, shining, very finely and faintly punctured; humeral angles rounding; apex obtuse; a crescent-shaped row of yellow dots connecting the humeral angles, these dots joined together near shoulders to make a short stripe; behind this row a transverse area filled with very small dots which in the same manner join to form a stripe at the margin; apex concolorous yellow-brown. These dots and fascia are bordered by bands of dark brown and are not symmetrically arranged bilaterally. Tegmina very dark brown, opaque, more than half covered by the pronotum, border wide and wrinkled. Legs and undersurface of body yellow. Tarsi and tips of tibiae fuscous-brown.

Type.—Female.

Length 3.6 mm.; width 2.4 mm.

Locality.—Bolivia.

The fore-wings of the genus *Tragopa* are most peculiar and are entirely different from the wings of other Membracidae. The wing of *T. decorata* figured is a good example of this interesting wing structure and is remarkable not only for the venation but also for the broad, corrugated limbus.