

for few days; rash during the second week; tympanites slight during early days; coated tongue; temperature 103° and slightly above for first few days, after that gradual decline until normal on fourteenth day; stools deodorized; no bad symptoms.

Treatment as in Case 2.

CASE 4. Miss G.B., æt. 19. Headache; coated tongue; rash appeared on fourth day; slight epistaxis at first; bowels constipated; delirium slight at night at commencement; temperature on first day, 103.2° , ranging a few points above 102° during first week, declining gradually until normal on ninth day; no tympanites.

Treatment as in Case 2.

CASE 5. Thomas R., æt. 28. Temperature at first visit, 103.1° , gradual fall until normal on tenth day; violent headache; no delirium; coated tongue throughout; epistaxis on several occasions; no rash; no tympanites; dilated pupils at first; enlargement of spleen; stools deodorized; lost flesh considerably during attack.

Treatment as in preceding case.

CASE 6. George G., æt. 28. Temperature 104.1° at first visit, between 103° and 104° for first week, gradual decline after that until normal on fourteenth day; delirium on first night; headache intense; constipation; rash during first week; nose bled frequently; no tympanites; stools lost offensive character after a few days; spleen enlarged; second slight rise of temperature occurred, lasting for three days, associated with pain and tenderness over the liver; phlebitis in left leg during convalescence.

Treatment as in other cases.

CASE 7. Helen B., æt. 5. Temperature 103.1° at first visit; child was at school three days before; temperature between 102° and 103° for first week, after that declined gradually until normal on evening of seventeenth day; no subsequent rise; headache intense; nose bled several times; tympanites at first, but disappeared; constipation obstinate at first, so that there was difficulty in securing elimination; stools lost offensive character; rash on second week, but scanty.

Treatment as in other cases, and in addition salines.

CASE 8. James M., æt. 26. Temperature in axilla at first visit 101° , continuing between 101° and 102° degrees for first week, then declining gradually until normal on twelfth day; coated tongue; pulpy gums; intense headache at first; disturbed sleep, but no delirium; widely-dilated pupils at commencement; spleen much enlarged and tender; flushed face; anorexia; no tympanites; stools lost offensive odor; rash, a few spots, during end of first week.

Treatment as in other cases, with daily enemata of glycerine to facilitate elimination.