

the violent movements accompanying the cough. To administer these tablets in the above conditions, place one tablet in the mouth, allowing it to dissolve slowly, swallowing the saliva. Exhibited in the grinding pains which precede and follow labor; in the uterine contractions which often lead to abortion; in the various neuralgias, and in all neuroses due to irregularities of menstruation, this combination affords immediate relief. In these last conditions, always instruct that tablets be crushed before taking.

EVERY PHYSICIAN KNOWS.

Every physician knows full well the advantages to be derived from the use of antikamnia in very many diseases, but a number of them are still lacking a knowledge of the fact that antikamnia in combination with various remedies, has a peculiarly happy effect; particularly is this the case when combined with salol. Salol is a most valuable remedy in many affections; and its usefulness seems to be enhanced by combining it with antikamnia. The rheumatoid conditions so often seen in various manifestations are wonderfully relieved by the use of this combination. After fevers, inflammations, etc., there frequently remain various painful and annoying conditions which may continue, namely: The severe headaches which occur after meningitis, a "stitch in the side" following pleurisy, the precordial pain of pericarditis and the painful stiffness of the joints which remain after a rheumatic attack—all these conditions are relieved by this combination called "Antikamnia & Salol Tablets" containing $2\frac{1}{2}$ grs., each of antikamnia and of salol and the dose of which is one or two every two or three hours. They are also recommended highly in the treatment of cases of both acute and chronic cystitis. The pain and burning is relieved to a marked degree. Salol neutralizes the uric acid and clears up the urine. This remedy is a reliable one in the treatment of diarrhoea, entero colitis, dysentery, etc. In dysentery, where there are bloody, slimy discharges, with tormina and tenesmus, a good dose of sulphate of magnesia, followed by two antikamnia and salol tablets every three hours will give results that are gratifying.

DR. HOWARD ON THE TREATMENT OF MALARIA.

"Since 1899 it has been my privilege to act as Medical Officer to the Universities' Mission to Central Africa. In that capacity I have had charge of a number of Europeans living in typical outlying stations in the neighborhood of Lake Nyasa, and during this period I have personally made observations and directed anti-malarial measures at all stations, so that my conclusions are based on prolonged study and practical knowledge of the difficulties which have to be encountered.