

and hæmorrhage. Last August, 1905, had a slight hæmorrhage from the lungs; since that time has been coughing and expectorating. On January 1st, 1906, had a severe hæmorrhage from the lungs. Temperature was 102; pulse, 108.

Lungs—Softening in the left apex. Injected with 12 grains January 6th, 1906. Cough almost ceased; appetite was good. He was put on iodine gr. one half every four hours and Easton's formula. Examined February 8th, coughs a few times in the morning. *Lungs*—only slight roughening of the breathing in the supraclavicular region.

Three cavity cases have been treated by this method. They all died—two of septic pneumonia and the other of heart failure. These cases were all septic. Both the streptococcus and tubercle bacilli were found in the sputa. The second case, Mr. S., was treated with antistreptococcic serum with reduction of temperature at the end.

Case XIII.—The last case, Mrs. B., age 25, came under treatment August 4th 1905, with active disease, a cavity in right apex, and one in left upper lobe about the third rib. Temperature running up to 103, and pulse 96 to 108. Tubercular diarrhoea. She received several injections and always with benefit, but as there was streptococcic infection the disease progressed. After treatment for a time with iodoform, streptococci serum was used, and by November 1st, 1905, her temperature became normal and her diarrhoea stopped. It was noticed toward the last that a combination treatment gave the best results. In treating the above cases the dose of iodoform was too large, and too far apart, and the patients were not kept at rest.

I have two cavity cases at present under treatment. They are receiving small injections often, from seven to twelve days, and iodine gr. one half every four hours after the reaction is over. They are slowly progressing.

The feeding and the care of the patient are carried out after latest methods, by fresh eggs, fats and fresh air.

Tuberculosis of the Bowels.

Case XIV.—Mr. B., came under treatment November 1st, 1905, with cough, purulent expectoration, fever and diarrhoea. Patient had a tubercular diarrhoea for two years.

Lungs—Cavity in right apex. Patient was very weak and having between five and seven stools a day.

He has been injected with 8 grains every twelve days, and iodine gr. one half every four hours after the reaction is over. His diarrhoea