

isms, which, by their accumulation, modify the soil on which the microbes develop. In their experiments made on rabbits, the investigators observed that any nutritive substance which had served as a culture medium for the pneumococci, even if it had been separated from the microbes by filtration, conferred on the animal immunity against the pneumonic infection. They next proved experimentally that the blood serum of a rabbit "vaccinated" against the pneumococcus may cure an animal infected with pneumonia. An intravenous injection of eight cubic centimetres of serum of an animal rendered refractory, practised twenty-four hours after the infection, produces a gradual fall in the febrile temperature, and hastens the recovery of the animal. In another series of researches, devoted to the study of the cause of the remedial action of the serum of inoculated animals, the same observers found that the pneumococcus, when introduced into the body of an animal, gives rise to the production of a "pneumotoxine," which may be isolated. This pneumotoxine produces a febrile reaction of several days' duration, after which they have noted in the fluids of the animal another substance, "anti-pneumotoxine," which has the power of neutralizing pneumotoxine. The blood serum of an animal on which immunity has been conferred contains anti-pneumotoxine, and it is this which seems to forward the recovery from the pneumonic infection. In the blood serum of patients affected with croupous pneumonia, they have also found pneumotoxine and anti-pneumotoxine, the former chiefly during the febrile period of the disease, the latter after the crisis. They also claim to have treated successfully rabbits suffering from pneumonia by injecting into these animals blood serum taken from a pneumonic patient after the crisis. Being assured by experiments made on themselves that man may support with impunity, and without any local and general reaction, injections of the serum of animals rendered refractory to Fraenkel's pneumococcus, the investigators treated six patients affected with pneumonia. Although the number treated was small, the result has been very encouraging. In fact, in all these patients a hypodermic injection of from four to six cubic centimetres of serum was followed at the end

of from six to twelve hours by a considerable fall in the temperature, with slowing of the pulse and respiration. These observations are especially noteworthy as confirming those made by Emmerich and Fowitzky, who claim that they have conferred immunity on the rabbit by means of hypodermic injections of attenuated cultures of the pneumococcus; but this immunity, they say, is incomplete. On rabbits infected by pneumococci, on the other hand, full immunity is obtained by intravenous injections of a culture having its entire virulence, but largely diluted. The liquid obtained by crushing the organs of an animal thus rendered refractory exercises on the pneumonic infection a sure remedial action when it is injected under the skin or into the abdominal cavity, and especially when it is thrown into the veins of the infected animal.—*Lancet*.

THE HEALTH OF VETERANS, OR TWENTY-FIVE YEARS AFTER THE WAR.—Dr. John L. Billings, of the army, has contributed to *The Forum* for January a brief study of the health of the survivors of the war, as judged by data compiled under the eleventh census. The author's intention is to show with regard to the troops of a single state, Massachusetts, what may at some future time be worked up for the Northern States as a whole. In that state about 40,000 veterans were reported as living on the first of June, 1890. These form about one-eighth part of the white male population over forty years old. Hence, if they were all equally healthy, the number of sickness cases reported by the census should be seven times as great among the latter as among the former. But the census indicates that there is four times as much sickness among veterans as among other males of the same age. Among the insane, however, the veterans furnished a much smaller proportion than the other males over forty years of age. The sickness statistics were especially high among the veterans from diarrhoeal diseases, rheumatism, and heart disease. This fact might have been anticipated, and may, in part at least, be set down as one of the entailments of service in the field. Dr. Billings infers that while the health of some men has been improved by their war discipline—even to the extent of the preservation of