

tively easy menstruations occurring recently, in spite of the displacement, were due entirely to Ergoapiol.

CASE 4.—Miss —, aged eighteen, had always been regular in menstruating. Could get no history of any previous disorder within patient's knowledge. Contracted a heavy cold about time of menstrual epoch, and was much alarmed by non-appearance of flow. Discomfort was not marked. Ergoapiol (Smith), one capsule three times a day, was prescribed. Reported later that flow was established in twenty-four hours after treatment was commenced. The delay in this case was about four days.

CASE 5.—Mrs. — consulted me, giving the following history: Three months previously had had a profuse uterine hemorrhage occurring about the time of menstrual period; as she had for a number of years menstruated only at intervals of about six or seven weeks, the fact that menstruation had been suspended for six weeks before the date of trouble was not especially significant. The hemorrhage, which was at no time alarming, had continued for several days. Since that time there had been an almost constant wasting and at times a considerable flow. Her condition was practically invalid. Examination revealed a gaping os, a cervix exceedingly tender and abraded, and a large uterus. Before resorting to curettement it seemed advisable to try other measures. Ergoapial (Smith), one capsule every three hours, was prescribed. In about twenty-four hours there was a decided increase in the discharge, which consisted of clots and considerable debris. There were some pains, of a cramp-like nature. The discharge began to grow less in about four days and ceased entirely in one week. There was a marked improvement in general condition. Local treatment entirely removed the tenderness and abraded condition of cervix. Ergoapol (Smith) was administered several days before next menstrual period and resulted in a very satisfactory period. In this case it appears to me the remedy saved the patient the ordeal of curettement, acting as a prompt uterine stimulant. Her condition locally and generally has since steadily improved.

---

To prevent the nausea following the administration of the various opium preparations, and even of the drug itself, add cannabis indica to the dose, and follow by small and repeated doses until danger of nausea has passed.—*Medical Council.*