

foundland and Labrador to Great Slave Lake and southward to Nova Scotia, northern New England, Quebec, Ontario and northern Michigan. I have seen but nine specimens from the following localities:

Hopedale, Labrador, Aug., 1917, 1 ♂, (W. W. Perrett) Ashwanipi River, Northern Quebec, July 25, 1917, 1 ♀, (Carnegie Museum Coll.); Isle d'Orleans, Que., Aug. 27, 1904, 1 ♂, (Walker); Algonquin Park, Ont., July 15, 1900, 1 ♂, (J. Macoun); De Grassi Pt., Ont., June 26, 1918, 1 ♂, (Walker); Profile Lake, N. H., July 20, 1918, 1 ♂, (R. H. Howe); Manistiqua River, Schoolcraft Co., Mich., 2 ♂'s, (A. F. Combs); Marquette, Mich., 1 ♀, (Coll. Hubbard & Schwarz, U. S. N. M.).

Somatochlora kennedyi, n. sp.

Male.—Occiput, frontal vesicle and upper part of frons metallic blue-black or green-black; sides of frons and a moderately broad, lower margin yellowish brown; postclypeus dark brown or black in centre, lateral lobes yellow; labrum black or deep brown, anteclypeus and labium pale yellow. Pile somewhat dense, blackish on top of head, pale yellowish on face. Rear of head black with whitish pile.

Prothorax greyish brown, anterior lobe edged with whitish, posterior lobe testaceous. Meso- and metathorax metallic gold-green with blue reflexions, the actual brilliancy of the colour much obscured by dense, light brownish pile; the following parts ochreous but not forming well-defined nor conspicuous spots:—the antealar sinus, the ventral edges of the mesepisternum, a narrow area below bases of wings, an oblong or ovate spot on the mesepimerum connected with the above area, an ill-defined blotch enclosing the metastigma, and the posterior half or more of the metepimerum. Fore coxæ black in front, dull yellowish above and behind, this colour extending over trochanters to outer surface of fore femora in their basal half.

Wings hyaline or tinged with yellow, especially along costa, sometimes entirely suffused with amber yellow; costa yellowish proximally, darkening beyond nodus; pterostigmata dark yellowish brown; hind wings with a yellow basal spot of variable size and