

$b\rho z' \quad q \quad \Delta S \quad \Delta a \Gamma \lesssim z.$

$\triangle A \triangle^{\circ} \Gamma \Gamma \Delta \delta \cdot \nabla \Lambda \rho' \cdot \triangleleft \triangle \Lambda \alpha^{\circ} \rho^{\circ} \Delta^{\circ} \text{ L} \Gamma \Delta \delta \cdot \nabla -$
 $\Lambda \rho' \Delta^{\circ} \text{ b} \Delta \delta \rho \rho', \text{)} (\alpha \text{ } \zeta^{\circ} \Delta^{\circ} \nabla \rho \alpha \delta \sigma \text{ b} \alpha \text{ b} \triangleleft \Delta^{\circ}$
 $\cdot \nabla \sigma \delta \delta \alpha, \text{ D} \text{ b} \Lambda \text{ L} \Gamma \triangle^{\circ} \Delta \sigma^{\circ} \text{ D} \rho \text{ i} \text{ b}^{\circ} \text{ b}^{\circ} \text{ }^{\times}$

$\sigma \cdot \Delta^{\dagger}(\rho) = \sigma_L \Gamma \Delta \Gamma^{\dagger} q \Delta \rho$ ከ $\sigma \cdot \langle \Gamma^{\dagger} \Delta \rangle$ ጋር
 $\sigma \otimes \rho \cdot b \cdot \langle \Delta^{\dagger} \rangle^*$

b7C σΛΓΔΣ∇ΛΓ'Δα, b6 `bΓΔΛΔL'ΔΣ) bPa
σbP<ΠΓ'Δα)*

[illegible][illegible][illegible]

$\triangleright \nabla \triangleleft$, $L_0 < \omega^2 \Delta t^2$, $\Delta L \nabla C$ $P \cap \langle d\sigma_9 - \Delta\sigma_2$; $b:\Delta$ ΔL $P\sigma^y b \cap r:\Delta\sigma_2$, $d\sigma_L$ $P \dot{P} \dot{P} \dot{Q} \dot{L}^\circ \times$