INDEX TO CONTENTS.

P	AGR.
Some Historical and Other Data on Circumcision	181
Treatment of Tapeworm by use of Morphine	186
Some of the Commoner Reflex Symptoms from Disease of the Rectum	
The Human Foot	i 90
Mechanics of the Foot	
Anaemia and its Treatment	
A Case of Septicemia Post-abortum	
Bubonic Plague in 1141 B.C	
Ichthyol in the Treatment of Corneal Ulcers	
Therapeutic Notes	
Editorial	

A STRONG STATEMENT A STRONG SOURCE

"In mild forms of syphilis with little or no glandular involvement, and in mixed sores, I immediately resort to mercauro, and I have never seen its equal as a remedial agent in syphilis. It is pre-eminently a new therapeutic agent, the physiological effect being subjudice. In latent lesions it is positively the best remedy. By abundant experience I am settled in my conviction as to its value, and more particularly so, when the iodides are not tolerated."

NEW YORK POLYCLINIC.—April 1897. SYPHILIS AND SEXUAL NEURASTHENIA:

By Herman F. Nordeman, M. D.,

Adjunct-Professor, Genito-Urinary Diseases, at the New York Polyclinic Surgeon to New York Surgical and Genito-Urinary Hospital, etc., etc.